

**ND HHS measles outbreak vaccination recommendations for children who live in or visit counties with ongoing measles transmission<sup>1</sup>**

Age	Number of previous vaccine doses	MMR vaccine recommendations
<b>0 – 6 months</b>	0	<ul style="list-style-type: none"> <li>• Vaccination is not recommended</li> </ul>
<b>6 – 11 months</b>	0	<ul style="list-style-type: none"> <li>• Should receive an early dose of MMR<sup>2</sup> vaccine immediately</li> <li>• Should receive two additional doses of MMR vaccine on the regular schedule               <ul style="list-style-type: none"> <li>○ First dose at 12-15 months</li> <li>○ Second dose at 4-6 years</li> </ul> </li> <li>• Receive each dose at least 28 days apart</li> </ul>
<b>1 – 17 years</b>	0	<ul style="list-style-type: none"> <li>• Should receive first dose immediately</li> <li>• Should receive second dose at least 28 days later</li> </ul>
	1	<ul style="list-style-type: none"> <li>• Should receive a second dose of MMR vaccine at least 28 days after first dose</li> </ul>
	2	<ul style="list-style-type: none"> <li>• Fully vaccinated; no additional doses needed</li> </ul>

<sup>1</sup> As of 4/16/2026 counties with ongoing measles transmission include: Grand Forks County, Ransom County

<sup>2</sup> Measles, Mumps, Rubella, and Varicella (MMRV) vaccine should not be utilized in infants under 12 months of age.

**ND HHS measles outbreak vaccination recommendations for adults who live in or visit counties with ongoing measles transmission<sup>1</sup>**

If you were born	Number of previous vaccine doses	MMR vaccine recommendations
<b>Before 1957</b>	N/A	<ul style="list-style-type: none"> <li>Likely exposed to measles as a child; vaccination not needed</li> </ul>
<b>Between 1957-1968</b>	0	<ul style="list-style-type: none"> <li>Should receive first dose of live-attenuated MMR vaccine immediately</li> </ul>
	1 dose of inactivated MMR vaccine	<ul style="list-style-type: none"> <li>Should receive second dose at least 28 days later</li> </ul>
	1 dose of live-attenuated MMR vaccine	<ul style="list-style-type: none"> <li>Should receive second dose</li> </ul>
<b>After 1968</b>	0	<ul style="list-style-type: none"> <li>Should receive first dose immediately</li> <li>Should receive second dose at least 28 days later</li> </ul>
	1	<ul style="list-style-type: none"> <li>Should receive a second dose of MMR vaccine at least 28 days after first dose</li> </ul>
	2	<ul style="list-style-type: none"> <li>Fully vaccinated; no additional doses needed</li> </ul>

<sup>1</sup> As of 4/16/2026 counties with ongoing measles transmission include: Grand Forks County, Ransom County

Last updated: 4/16/2026

## Measles, Mumps, and Rubella (MMR) Vaccine Recommendations for Specific Populations

<b>Pregnant Women</b>	<ul style="list-style-type: none"><li>• Vaccination is not recommended during pregnancy</li></ul>
<b>Severely Immunocompromised individuals</b>	<ul style="list-style-type: none"><li>• MMR vaccine is not recommended for individuals with severe immunodeficiency</li><li>• Severe immunodeficiency includes hematologic and solid tumors, receipt of chemotherapy, congenital immunodeficiency, long-term immunosuppressive therapy or patients with HIV infection who are severely immunocompromised.</li></ul>
<b>Healthcare Personnel</b>	<ul style="list-style-type: none"><li>• Healthcare personnel without presumptive evidence of immunity should get two doses of MMR vaccine, at least 28 days apart.</li></ul>