NORTH DAKOTA

2011 Land Line Only Module Variables Report

Behavioral Risk Factor Surveillance System

July 26, 2012





MODULE 2: DIABETES CONTENTS

	EN THE DENOMINATOR IS NOT BASED ON THE TOTAL POPULATION SURVEYED, THE TITLE CONTAINS A DESCRIPTION OF WHAT POPULAT EXCLUDED FROM THE DENOMINATOR. IF THE TITLE DOES NOT DESCRIBE THE DENOMINATOR, THE TABLE IS BASED ON ALL RESPONDEN TO THE STATES SURVEY.	
* * *	*******************************	* * *
TABLE		PAGE
	HOW OLD WERE YOU WHEN YOU WERE TOLD YOU HAVE DIABETES? (DIABAGE2)	
1	MALES AND FEMALES	4
2	MALES ONLY	5
3	FEMALES ONLY	6
	ARE YOU NOW TAKING INSULIN? (INSULIN)	
4	MALES AND FEMALES	7
5	MALES ONLY	8
6	FEMALES ONLY	9
	ABOUT HOW OFTEN DO YOU CHECK YOUR BLOOD FOR GLUCOSE OR SUGAR? INCLUDE TIMES WHEN CHECKED BY A FAMILY MEMBER OR FRIEND, BUT DO NOT INCLUDE TIMES WHEN CHECKED BY A HEALTH PROFESSIONAL. (BLDSUGAR)	
7	MALES AND FEMALES	10
8	MALES ONLY	11
9	FEMALES ONLY	12
	ABOUT HOW OFTEN DO YOU CHECK YOUR FEET FOR ANY SORES OR IRRITATIONS? INCLUDE TIMES WHEN CHECKED BY A FAMILY MEMBER OR FRIEND, BUT DO NOT INCLUDE TIMES WHEN CHECKED BY A HEALTH PROFESSIONAL. (FEETCHK2)	
10	MALES AND FEMALES	13
11	MALES ONLY	14
12	FEMALES ONLY	15
	ABOUT HOW MANY TIMES IN THE PAST 12 MONTHS HAVE YOU SEEN A DOCTOR, NURSE, OR OTHER HEALTH PROFESSIONAL FOR YOUR DIABETES? (DOCTDIAB)	
13	MALES AND FEMALES	16
14	MALES ONLY	17
15	FEMALES ONLY	18
	A TEST FOR "A ONE C" MEASURES THE AVERAGE LEVEL OF BLOOD SUGAR OVER THE PAST THREE MONTHS. ABOUT HOW MANY TIMES IN THE PAST 12 MONTHS HAS A DOCTOR, NURSE, OR OTHER HEALTH PROFESSIONAL CHECKED YOU FOR "A ONE C"? (CHKHEMO3)	
16	MALES AND FEMALES	19
17	MALES ONLY	20
18	FEMALES ONLY	21

MODULE 2: DIABETES CONTENTS

NOT		TTO L
	EN THE DENOMINATOR IS NOT BASED ON THE TOTAL POPULATION SURVEYED, THE TITLE CONTAINS A DESCRIPTION OF WHAT POPULA	
IS	EXCLUDED FROM THE DENOMINATOR. IF THE TITLE DOES NOT DESCRIBE THE DENOMINATOR, THE TABLE IS BASED ON ALL RESPONDE	NTS
also also also a	TO THE STATES SURVEY.	ale de de de
***	**************************************	* * * *
TABLE		PAGE
	ABOUT HOW MANY TIMES IN THE PAST 12 MONTHS HAS A HEALTH PROFESSIONAL CHECKED YOUR FEET FOR ANY SORES OR IRRITATIONS? (FEETCHK)	
19	MALES AND FEMALES	22
20	MALES ONLY	23
21	FEMALES ONLY	24
	WHEN WAS THE LAST TIME YOU HAD AN EYE EXAM IN WHICH THE PUPILS WERE DILATED? THIS WOULD HAVE MADE YOU TEMPORARILY SENSITIVE TO BRIGHT LIGHT. (EYEEXAM)	
22	MALES AND FEMALES	25
23	MALES ONLY	26
24	FEMALES ONLY	27
	HAS A DOCTOR EVER TOLD YOU THAT DIABETES HAS AFFECTED YOUR EYES OR THAT YOU HAD RETINOPATHY? (DIABEYE)	
25	MALES AND FEMALES	28
26	MALES ONLY	29
27	FEMALES ONLY	30
	HAVE YOU EVER TAKEN A COURSE OR CLASS IN HOW TO MANAGE YOUR DIABETES YOURSELF? (DIABEDU)	
28	MALES AND FEMALES	31
29	MALES ONLY	32
30	FEMALES ONLY	33

MODULE 2: DIABETES

TABLE 1: HOW OLD WERE YOU WHEN YOU WERE TOLD YOU HAVE DIABETES? (DIABAGE2)

Denominator excludes: Respondents with do not know/refused/missing responses Denominator excludes: Respondents that reported not being diagnosed with diabetes

Denominator excludes: Women respondents that reported being diagnosed with diabetes only during pregnancy

Data was weighted using _LANDWT.

DEMOGRAPHIC	RESPONDE	NT NUMBER	1-	15 yea	rs old	16	-30 yea	ars old	31	-45 yea	ars old	46	-60 yea	ars old	6	1+ year	rs old
GROUPS	TOTAL	WEIGHTED	N	%	C.I. (95%)	N	%	C.I. (95%)	N	%	C.I. (95%)	N	%	C.I. (95%)	N	%	C.I. (95%)
TOTAL	477	38,881	7	2.6	00.4-04.8	23	7.7	04.2-11.2	89	23.5	18.4-28.6	211	41.5	36.2-46.8	147	24.8	20.7-28.9
Male	213	20,370	3	2.3	00.0-05.0	10	6.2	01.9-10.5	39	23.9	16.5-31.3	99	42.2	34.8-49.6	62	25.4	19.1-31.7
Female	264	18,511	4	2.9	00.0-06.2	13	9.3	03.8-14.8	50	23.0	15.7-30.3	112	40.7	33.6-47.8	85	24.1	18.8-29.4
Non-Native	444	34,938	6	2.4	00.2-04.6	19	7.6	03.9-11.3	81	22.1	17.0-27.2	198	41.8	36.5-47.1	140	26.1	21.6-30.6
45-54	63	7,678	2	2.6	00.0-06.7	7	11.6	03.0-20.2	32	57.2	43.7-70.7	22	28.6	16.8-40.4			
55-64	135	10,843	2	1.4	00.0-03.4	4	2.7	00.0-05.4	28	22.9	13.1-32.7	93	66.4	56.2-76.6	8	6.6	01.1-12.1
65+	262	17,188				5	2.0	00.2-03.8	23	7.3	04.2-10.4	96	39.2	32.3-46.1	138	51.5	44.6-58.4
H.S. or G.E.D.	171	12,534	2	3.0	00.0-07.1	6	5.4	00.3-10.5	40	29.9	20.7-39.1	68	37.7	29.3-46.1	55	24.0	17.3-30.7
Some Post-H.S.	149	14,606	4	3.1	00.0-06.8	12	13.2	05.8-20.6	22	19.8	11.4-28.2	78	46.1	36.9-55.3	33	17.8	11.5-24.1
College Graduate	107	6,230	1	2.9	00.0-08.6	3	2.4	00.0-05.1	24	26.7	15.7-37.7	51	48.3	36.9-59.7	28	19.7	12.1-27.3
Less than \$15,000	57	5,540	2	3.7	00.0-09.2	4	7.9	00.1-15.7	8	26.4	08.6-44.2	25	36.0	21.5-50.5	18	26.1	13.2-39.0
\$15,000- 24,999	88	7,033	2	6.2	00.0-14.4	5	10.9	00.5-21.3	13	16.8	06.2-27.4	36	35.9	23.9-47.9	32	30.2	19.0-41.4
\$25,000- 34,999	69	5,246				2	4.6	00.0-11.5	15	23.5	10.4-36.6	28	39.3	25.8-52.8	24	32.6	20.3-44.9
\$35,000- 49,999	64	5,164	2	2.8	00.0-06.9	3	7.0	00.0-16.8	10	21.4	07.7-35.1	29	43.3	29.4-57.2	20	25.4	14.2-36.6
\$50,000- 74,999	58	5,407	1	4.2	00.0-12.0	3	11.1	00.0-23.1	15	30.5	15.4-45.6	28	42.6	27.9-57.3	11	11.6	04.3-18.9
\$75,000+	57	4,546				5	11.2	01.2-21.2	16	34.6	19.1-50.1	25	40.0	25.5-54.5	11	14.2	05.2-23.2

NOTE:

^{&#}x27;% = Percentage, C.I.(95%) = Confidence Interval (at 95 percent probability level). Percentages are weighted to population characteristics. Rows with less than 50 observations have been suppressed.

MODULE 2: DIABETES

TABLE 2: HOW OLD WERE YOU WHEN YOU WERE TOLD YOU HAVE DIABETES? (DIABAGE2)

Denominator excludes: Respondents who are female or respondents with do not know/refused or with missing responses Denominator excludes: Respondents that reported not being diagnosed with diabetes

Denominator excludes: Women respondents that reported being diagnosed with diabetes only during pregnancy

Data was weighted using _LANDWT.

DEMOGRAPHIC	RESPONDE	NT NUMBER	1	-15 yea	rs old	16	-30 yea	ars old	31	-45 yea	ars old	46	6-60 yea	rs old	6	1+ yea	rs old
GROUPS	TOTAL	WEIGHTED	N	%	C.I. (95%)	N	%	C.I. (95%)	N	%	C.I. (95%)	N	%	C.I. (95%)	N	%	C.I. (95%)
Male	213	20,370	3	2.3	00.0-05.0	10	6.2	01.9-10.5	39	23.9	16.5-31.3	99	42.2	34.6-49.8	62	25.4	19.1-31.7
Non-Native	198	18,839	3	2.5	00.0-05.4	8	5.7	01.4-10.0	36	24.4	16.6-32.2	93	41.9	34.1-49.7	58	25.5	18.8-32.2
55-64	68	5,946	1	1.6	00.0-04.7	3	3.8	00.0-08.5	12	20.4	08.2-32.6	46	64.8	51.1-78.5	6	9.4	00.4-18.4
65+	110	8,897				3	2.6	00.0-05.7	7	5.2	01.1-09.3	44	40.4	30.2-50.6	56	51.9	41.7-62.1
H.S. or G.E.D.	65	5,936	2	6.3	00.0-14.9	2	5.5	00.0-14.1	15	28.5	15.2-41.8	26	35.5	22.8-48.2	20	24.2	13.8-34.6
Some Post-H.S.	61	6,884	1	1.4	00.0-04.1	4	8.9	00.1-17.7	7	21.9	07.6-36.2	37	49.7	35.4-64.0	12	18.2	07.8-28.6
College Graduate	57	3,810				2	2.4	00.0-05.7	15	33.4	17.7-49.1	27	46.1	30.8-61.4	13	18.1	07.9-28.3

^{&#}x27;% = Percentage, C.I.(95%) = Confidence Interval (at 95 percent probability level). Percentages are weighted to population characteristics. Rows with less than 50 observations have been suppressed.

MODULE 2: DIABETES

TABLE 3: HOW OLD WERE YOU WHEN YOU WERE TOLD YOU HAVE DIABETES? (DIABAGE2)

Denominator excludes: Respondents who are male or respondents with do not know/refused or with missing responses Denominator excludes: Respondents that reported not being diagnosed with diabetes

Denominator excludes: Women respondents that reported being diagnosed with diabetes only during pregnancy

Data was weighted using _LANDWT.

DEMOGRAPHIC	RESPONDE	NT NUMBER	1	-15 yea	rs old	16	-30 yea	ars old	31	-45 yea	ars old	46	-60 yea	ars old	6	1+ yea	rs old
GROUPS	TOTAL	WEIGHTED	N	%	C.I. (95%)	N	%	C.I. (95%)	N	%	C.I. (95%)	N	%	C.I. (95%)	N	%	C.I. (95%)
Female	264	18,511	4	2.9	00.0-06.2	13	9.3	03.8-14.8	50	23.0	15.7-30.3	112	40.7	33.6-47.8	85	24.1	18.8-29.4
Non-Native	246	16,099	3	2.3	00.0-05.4	11	9.9	03.8-16.0	45	19.4	13.5-25.3	105	41.7	34.4-49.0	82	26.8	20.9-32.7
55-64	67	4,898	1	1.1	00.0-03.3	1	1.3	00.0-03.8	16	26.1	10.4-41.8	47	68.3	52.6-84.0	2	3.3	00.0-07.8
65+	152	8,291				2	1.4	00.0-03.4	16	9.6	04.9-14.3	52	37.9	29.1-46.7	82	51.2	42.4-60.0
H.S. or G.E.D.	106	6,598				4	5.3	00.0-11.0	25	31.2	18.7-43.7	42	39.6	28.4-50.8	35	23.9	15.7-32.1
Some Post-H.S.	88	7,721	3	4.7	00.0-11.2	8	17.0	05.6-28.4	15	17.9	08.7-27.1	41	42.9	30.9-54.9	21	17.5	09.9-25.1
College Graduate	50	2,420	1	7.5	00.0-21.4	1	2.6	00.0-07.5	9	16.1	05.5-26.7	24	51.6	35.1-68.1	15	22.2	10.8-33.6
\$15,000- 24,999	59	4,069	2	10.8	00.0-24.5	3	14.6	00.0-31.1	11	19.1	06.8-31.4	21	29.5	15.6-43.4	22	26.1	13.9-38.3

^{&#}x27;% = Percentage, C.I.(95%) = Confidence Interval (at 95 percent probability level). Percentages are weighted to population characteristics. Rows with less than 50 observations have been suppressed.

TABLE 4: ARE YOU NOW TAKING INSULIN? (INSULIN)

Denominator excludes: Respondents with do not know/refused/missing responses
Denominator excludes: Respondents that reported not being diagnosed with diabetes
Denominator excludes: Women respondents that reported being diagnosed with diabetes only during pregnancy

Data was weighted using LANDWT.

	Data w	as weighte	a usin	.9 _ьа	NDWI.		No N % C.I.								
DEMOGRAPHIC	RESPONDE	NT NUMBER		Yes	.		No								
GROUPS	TOTAL	WEIGHTED	N	%	C.I. (95%)	N	%	C.I. (95%)							
TOTAL	503	41,116	172	35.2	30.1-40.3	331	64.8	59.7-69.9							
Male	219	20,801	77	36.0	28.6-43.4	142	64.0	56.6-71.4							
Female	284	20,314	95	34.3	27.6-41.0	189	65.7	59.0-72.4							
Non-Native	467	36,873	158	35.7	30.4-41.0	309	64.3	59.0-69.6							
45-54	63	7,678	21	35.0	21.1-48.9	42	65.0	51.1-78.9							
55-64	140	11,378	45	29.7	21.1-38.3	95	70.3	61.7-78.9							
65+	280	18,603	98	37.0	30.5-43.5	182	63.0	56.5-69.5							
Less Than H.S.	57	6,232	23	38.7	24.4-53.0	34	61.3	47.0-75.6							
H.S. or G.E.D.	180	13,394	59	34.8	26.4-43.2	121	65.2	56.8-73.6							
Some Post-H.S.	157	15,186	53	34.8	26.0-43.6	104	65.2	56.4-74.0							
College Graduate	107	6,230	35	32.6	22.2-43.0	72	67.4	57.0-77.8							
Less than \$15,000	62	5,903	26	36.1	22.2-50.0	36	63.9	50.0-77.8							
\$15,000- 24,999	93	7,386	32	35.9	23.6-48.2	61	64.1	51.8-76.4							
\$25,000- 34,999	76	5,967	20	24.2	13.2-35.2	56	75.8	64.8-86.8							
\$35,000-49,999	65	5,354	27	41.8	27.9-55.7	38	58.2	44.3-72.1							
\$50,000-74,999	58	5,329	18	42.4	26.5-58.3	40	57.6	41.7-73.5							
\$75,000+	57	4,546	16	28.5	14.2-42.8	41	71.5	57.2-85.8							

NOTE:

^{&#}x27;% = Percentage, C.I.(95%) = Confidence Interval (at 95 percent probability level). Percentages are weighted to population characteristics.

Rows with less than 50 observations have been suppressed.

TABLE 5: ARE YOU NOW TAKING INSULIN? (INSULIN)

Denominator excludes: Respondents who are female or respondents with do not know/refused or with missing responses

Denominator excludes: Respondents that reported not being diagnosed with diabetes

Denominator excludes: Women respondents that reported being diagnosed with diabetes only during pregnancy

Data was weighted using _LANDWT.

DEMOGRAPHIC	RESPONDE	NT NUMBER		Yes	3		No	
GROUPS	TOTAL	WEIGHTED	N	%	C.I. (95%)	N	%	C.I. (95%)
Male	219	20,801	77	36.0	28.4-43.6	142	64.0	56.4-71.6
Non-Native	203	19,213	73	37.3	29.3-45.3	130	62.7	54.7-70.7
55-64	69	5,999	22	27.4	16.2-38.6	47	72.6	61.4-83.8
65+	115	9,275	44	40.0	30.0-50.0	71	60.0	50.0-70.0
H.S. or G.E.D.	67	6,032	23	42.2	28.1-56.3	44	57.8	43.7-71.9
Some Post-H.S.	63	6,962	20	29.4	16.3-42.5	43	70.6	57.5-83.7
College Graduate	57	3.810	20	31.4	18.5-44.3	37	68.6	55.7-81.5

NOTE:

^{&#}x27;% = Percentage, C.I.(95%) = Confidence Interval (at 95 percent probability level). Percentages are weighted to population characteristics.

Rows with less than 50 observations have been suppressed.

TABLE 6: ARE YOU NOW TAKING INSULIN? (INSULIN)

Denominator excludes: Respondents who are male or respondents with do not know/refused or with missing responses

Denominator excludes: Respondents that reported not being diagnosed with diabetes

Denominator excludes: Women respondents that reported being diagnosed with diabetes only during pregnancy

Data was weighted using _LANDWT.

DEMOGRAPHIC	RESPONDE	NT NUMBER		Yes	5		No	
GROUPS	TOTAL	WEIGHTED	N	%	C.I. (95%)	N	%	C.I. (95%)
Female	284	20,314	95	34.3	27.4-41.2	189	65.7	58.8-72.6
Non-Native	264	17,660	85	34.0	27.1-40.9	179	66.0	59.1-72.9
55-64	71	5,379	23	32.4	19.3-45.5	48	67.6	54.5-80.7
65+	165	9,328	54	34.0	25.8-42.2	111	66.0	57.8-74.2
H.S. or G.E.D.	113	7,362	36	28.8	19.6-38.0	77	71.2	62.0-80.4
Some Post-H.S.	94	8,224	33	39.4	27.4-51.4	61	60.6	48.6-72.6
College Graduate	50	2,420	15	34.6	18.1-51.1	35	65.4	48.9-81.9
\$15,000- 24,999	63	4.379	21	37.6	21.5-53.7	42	62.4	46.3-78.5

^{&#}x27;% = Percentage, C.I.(95%) = Confidence Interval (at 95 percent probability level). Percentages are weighted to population characteristics.

Rows with less than 50 observations have been suppressed.

MODULE 2: DIABETES

TABLE 7: ABOUT HOW OFTEN DO YOU CHECK YOUR BLOOD FOR GLUCOSE OR SUGAR? INCLUDE TIMES WHEN CHECKED BY A FAMILY MEMBER OR FRIEND, BUT DO NOT INCLUDE TIMES WHEN CHECKED BY A HEALTH PROFESSIONAL. (BLDSUGAR)

> Denominator excludes: Respondents with do not know/refused/missing responses Denominator excludes: Respondents that reported not being diagnosed with diabetes Denominator excludes: Women respondents that reported being diagnosed with diabetes only during pregnancy

> > Data was weighted using _LANDWT.

DEMOGRAPHIC	RESPONDE	NT NUMBER		Neve	r	<	1 time p	er Day	1-	5 times p	per Day	> !	5 times p	er Day
GROUPS	TOTAL	WEIGHTED	N	%	C.I. (95%)	N	%	C.I. (95%)	N	%	C.I. (95%)	N	%	C.I. (95%)
TOTAL	501	40,975	52	11.2	07.7-14.7	121	23.7	19.4-28.0	321	63.2	58.1-68.3	7	1.9	00.3-03.5
Male	218	20,731	27	12.0	06.7-17.3	66	29.8	22.9-36.7	120	55.5	47.9-63.1	5	2.8	00.1-05.5
Female	283	20,244	25	10.4	05.5-15.3	55	17.4	12.5-22.3	201	71.2	64.7-77.7	2	1.1	00.0-02.9
Non-Native	465	36,733	50	12.0	08.1-15.9	117	25.2	20.7-29.7	291	60.6	55.3-65.9	7	2.2	00.4-04.0
45-54	63	7,678	7	13.4	03.4-23.4	13	17.3	07.3-27.3	40	64.7	51.4-78.0	3	4.6	00.0-10.1
55-64	138	11,251	17	13.6	06.5-20.7	37	29.4	20.4-38.4	83	56.1	46.3-65.9	1	8.0	00.0-02.4
65+	280	18,590	24	7.0	04.1-09.9	68	25.2	19.3-31.1	186	67.1	61.0-73.2	2	0.7	00.0-01.7
Less Than H.S.	56	6,120	4	7.5	00.0-16.9	17	28.2	15.3-41.1	35	64.3	49.8-78.8			
H.S. or G.E.D.	180	13,393	15	9.5	04.2-14.8	40	21.1	14.2-28.0	123	67.2	59.0-75.4	2	2.2	00.0-05.5
Some Post-H.S.	156	15,159	17	11.2	05.1-17.3	36	23.2	15.8-30.6	99	62.6	53.8-71.4	4	2.9	00.0-06.0
College Graduate	107	6,230	16	18.4	08.0-28.8	28	26.1	16.9-35.3	62	54.7	43.3-66.1	1	8.0	00.0-02.4
Less than \$15,000	62	5,903	6	8.3	00.9-15.7	10	15.3	05.3-25.3	44	72.4	59.5-85.3	2	4.1	00.0-10.2
\$15,000- 24,999	93	7,307	7	11.4	00.2-22.6	25	24.2	14.0-34.4	61	64.5	51.8-77.2			
\$25,000- 34,999	76	5,967	9	14.4	03.6-25.2	16	23.2	11.6-34.8	51	62.4	49.1-75.7			
\$35,000- 49,999	64	5,260	4	6.1	00.0-12.6	13	22.2	10.0-34.4	45	68.9	55.8-82.0	2	2.8	00.0-06.7
\$50,000- 74,999	58	5,430	5	8.8	00.8-16.8	21	32.4	19.1-45.7	30	52.3	37.0-67.6	2	6.6	00.0-15.6
\$75,000+	57	4,546	8	13.3	03.3-23.3	18	33.3	19.0-47.6	31	53.4	38.3-68.5			

NOTE:

^{&#}x27;% = Percentage, C.I.(95%) = Confidence Interval (at 95 percent probability level). Percentages are weighted to population characteristics. Rows with less than 50 observations have been suppressed.

MODULE 2: DIABETES

TABLE 8: ABOUT HOW OFTEN DO YOU CHECK YOUR BLOOD FOR GLUCOSE OR SUGAR? INCLUDE TIMES WHEN CHECKED BY A FAMILY MEMBER OR FRIEND, BUT DO NOT INCLUDE TIMES WHEN CHECKED BY A HEALTH PROFESSIONAL. (BLDSUGAR)

Denominator excludes: Respondents who are female or respondents with do not know/refused or with missing responses

Denominator excludes: Respondents that reported not being diagnosed with diabetes

Denominator excludes: Women respondents that reported being diagnosed with diabetes only during pregnancy

Data was weighted using _LANDWT.

DEMOGRAPHIC	RESPONDE	NT NUMBER		Neve	r	<	1 time p	er Day	1-	5 times p	er Day	> !	5 times _l	oer Day											
GROUPS	TOTAL	WEIGHTED	N	%	C.I. (95%)	N	%	C.I. (95%)	N	%	C.I. (95%)	N	%	C.I. (95%)											
Male	218	20,731	27	12.0	06.7-17.3	66	29.8	22.9-36.7	120	55.5	47.9-63.1	5	2.8	00.1-05.5											
Non-Native	202	19,143	25	12.1	06.6-17.6	64	31.4	24.1-38.7	108	53.4	45.4-61.4	5	3.0	00.1-05.9											
55-64	68	5,905	10	16.8	05.4-28.2	24	38.4	25.3-51.5	33	43.2	30.1-56.3	1	1.6	00.0-04.7											
65+	115	9,298	12	7.8	03.1-12.5	31	28.3	18.9-37.7	70	62.5	52.7-72.3	2	1.3	00.0-03.3											
H.S. or G.E.D.	67	6,032	8	11.1	03.1-19.1	19	23.6	13.2-34.0	38	60.3	47.2-73.4	2	4.9	00.0-12.3											
Some Post-H.S.	63	7,003	6	7.4	00.9-13.9	19	33.0	19.9-46.1	36	56.4	42.5-70.3	2	3.2	00.0-07.7											
College Graduate	57	3,810	11	25.3	10.2-40.4	17	29.6	16.7-42.5	28	43.8	29.1-58.5	1	1.4	00.0-04.1											

^{&#}x27;% = Percentage, C.I.(95%) = Confidence Interval (at 95 percent probability level). Percentages are weighted to population characteristics.

Rows with less than 50 observations have been suppressed.

MODULE 2: DIABETES

TABLE 9: ABOUT HOW OFTEN DO YOU CHECK YOUR BLOOD FOR GLUCOSE OR SUGAR? INCLUDE TIMES WHEN CHECKED BY A FAMILY MEMBER OR FRIEND, BUT DO NOT INCLUDE TIMES WHEN CHECKED BY A HEALTH PROFESSIONAL. (BLDSUGAR)

> Denominator excludes: Respondents who are male or respondents with do not know/refused or with missing responses Denominator excludes: Respondents that reported not being diagnosed with diabetes Denominator excludes: Women respondents that reported being diagnosed with diabetes only during pregnancy

> > Data was weighted using LANDWT.

DEMOGRAPHIC	NT NUMBER		Neve	er	<	1 time p	er Day	1-	5 times p	per Day	> !	5 times _l	per Day	
GROUPS	TOTAL	WEIGHTED	N	%	C.I. (95%)	N	%	C.I. (95%)	N	%	C.I. (95%)	N	%	C.I. (95%)
Female	283	20,244	25	10.4	05.5-15.3	55	17.4	12.5-22.3	201	71.2	64.7-77.7	2	1.1	00.0-02.9
Non-Native	263	17,590	25	11.9	06.4-17.4	53	18.5	13.4-23.6	183	68.4	61.5-75.3	2	1.3	00.0-03.3
55-64 65+	70	5,345	7 12	10.1 6.1	02.3-17.9 02.4-09.8	13 37	19.5 22.1	08.5-30.5 15.2-29.0	50 116	70.4 71.8	57.7-83.1 64.4-79.2			
	165	9,292	12											
H.S. or G.E.D.	113	7,361	7	8.2	01.3-15.1	21	19.0	09.8-28.2	85	72.8	62.4-83.2			
Some Post-H.S.	93	8,155	11	14.5	04.9-24.1	17	14.8	07.5-22.1	63	67.9	56.5-79.3	2	2.7	00.0-06.8
College Graduate	50	2,420	5	7.5	00.4-14.6	11	20.6	08.6-32.6	34	72.0	58.5-85.5			
\$15,000- 24,999	64	4,412	6	12.2	00.0-26.3	14	17.8	08.0-27.6	44	70.0	54.9-85.1			

^{&#}x27;% = Percentage, C.I.(95%) = Confidence Interval (at 95 percent probability level). Percentages are weighted to population characteristics. Rows with less than 50 observations have been suppressed.

MODULE 2: DIABETES

TABLE 10: ABOUT HOW OFTEN DO YOU CHECK YOUR FEET FOR ANY SORES OR IRRITATIONS? INCLUDE TIMES WHEN CHECKED BY A FAMILY MEMBER OR FRIEND, BUT DO NOT INCLUDE TIMES WHEN CHECKED BY A HEALTH PROFESSIONAL. (FEETCHK2)

Denominator excludes: Respondents with do not know/refused/missing responses Denominator excludes: Respondents that reported not being diagnosed with diabetes Denominator excludes: Women respondents that reported being diagnosed with diabetes only during pregnancy

Data was weighted using _LANDWT.

DEMOGRAPHIC	RESPONDEN	NT NUMBER	Ti	imes pe	er Day	Tir	nes pe	r Week	Tin	nes per	Month	Tir	nes pe	r Year		Neve	er		No F	eet
GROUPS	TOTAL	WEIGHTED	N	%	C.I. (95%)	N	%	C.I. (95%)	N	%	C.I. (95%)	N	%	C.I. (95%)	N	%	C.I. (95%)	N	%	C.I. (95%)
TOTAL	494	40,080	281	56.6	51.3-61.9	98	18.8	14.7-22.9	33	7.5	04.4-10.6	5	1.1	00.1-02.1	73	15.3	11.6-19.0	4	0.8	00.0-01.6
Male	215	20,102	111	50.3	42.5-58.1	46	19.7	13.6-25.8	14	8.1	03.4-12.8	4	2.0	00.0-04.0	38	19.0	12.9-25.1	2	0.9	00.0-02.1
Female	279	19,978	170	62.9	55.8-70.0	52	17.8	12.3-23.3	19	6.9	02.8-11.0	1	0.2	00.0-00.6	35	11.5	07.2-15.8	2	0.7	00.0-01.7
Non-Native	458	35,838	255	54.8	49.3-60.3	91	18.3	14.2-22.4	33	8.4	04.9-11.9	5	1.2	00.0-02.4	70	16.4	12.3-20.5	4	0.9	00.1-01.7
45-54	63	7,678	35	55.1	41.0-69.2	13	19.4	08.6-30.2	7	11.7	01.5-21.9				8	13.9	04.3-23.5			
55-64	138	11,155	92	68.3	59.3-77.3	24	15.9	08.8-23.0	10	7.5	02.6-12.4	3	2.6	00.0-05.7	9	5.7	01.8-09.6			
65+	274	18,020	144	51.2	44.5-57.9	58	20.2	15.1-25.3	15	5.2	02.5-07.9	2	8.0	00.0-02.0	51	20.9	15.0-26.8	4	1.7	00.0-03.5
Less Than H.S.	52	5,625	28	57.9	42.6-73.2	7	10.0	02.4-17.6							17	32.1	17.6-46.6			
H.S. or G.E.D.	178	13,149	103	55.4	46.6-64.2	35	22.2	14.2-30.2	11	5.1	01.8-08.4	5	3.3	00.2-06.4	22	12.7	06.8-18.6	2	1.3	00.0-03.1
Some Post-H.S.	155	15,003	91	58.7	49.7-67.7	26	14.7	08.6-20.8	16	12.9	05.6-20.2				20	12.7	06.8-18.6	2	0.9	00.0-02.3
College Graduate	107	6,230	58	53.0	41.6-64.4	30	29.4	19.0-39.8	5	5.9	00.2-11.6				14	11.7	05.2-18.2			
Less than \$15,000	61	5,674	32	49.5	33.8-65.2	14	21.0	07.7-34.3	4	6.2	00.0-12.9				11	23.3	10.0-36.6			
\$15,000- 24,999	93	7,385	53	56.9	44.2-69.6	22	24.1	13.5-34.7	7	9.6	00.2-19.0	1	1.0	00.0-03.0	9	7.5	02.2-12.8	1	0.9	00.0-02.7
\$25,000- 34,999	76	5,967	49	64.2	51.7-76.7	8	9.2	02.5-15.9	3	3.4	00.0-07.5	1	1.3	00.0-03.8	14	20.6	09.4-31.8	1	1.3	00.0-03.8
\$35,000- 49,999	64	5,245	40	66.0	53.1-78.9	12	15.7	06.5-24.9	4	8.0	00.2-15.8	2	4.7	00.0-11.2	6	5.5	00.4-10.6			
\$50,000- 74,999	57	5,239	32	55.4	39.9-70.9	12	19.6	08.4-30.8	5	13.6	00.1-27.1				8	11.4	02.8-20.0			
\$75,000+	57	4,546	30	49.0	33.7-64.3	12	22.2	07.9-36.5	5	7.6	00.3-14.9				10	21.3	08.4-34.2			

NOTE:

^{&#}x27;% = Percentage, C.I.(95%) = Confidence Interval (at 95 percent probability level). Percentages are weighted to population characteristics. Rows with less than 50 observations have been suppressed.

MODULE 2: DIABETES

TABLE 11: ABOUT HOW OFTEN DO YOU CHECK YOUR FEET FOR ANY SORES OR IRRITATIONS? INCLUDE TIMES WHEN CHECKED BY A FAMILY MEMBER OR FRIEND, BUT DO NOT INCLUDE TIMES WHEN CHECKED BY A HEALTH PROFESSIONAL. (FEETCHK2)

Denominator excludes: Respondents who are female or respondents with do not know/refused or with missing responses

Denominator excludes: Respondents that reported not being diagnosed with diabetes

Denominator excludes: Women respondents that reported being diagnosed with diabetes only during pregnancy

Data was weighted using LANDWT.

DEMOGRAPHIC	RESPONDE	NT NUMBER	Т	imes pe	er Day	Tir	mes pe	r Week	Tin	nes pei	Month	Tir	nes pe	er Year		Nev	er		No F	eet
GROUPS	TOTAL	WEIGHTED	N	%	C.I. (95%)	N	%	C.I. (95%)	N	%	C.I. (95%)	N	%	C.I. (95%)	N	%	C.I. (95%)	N	%	C.I. (95%)
Male	215	20,102	111	50.3	42.5-58.1	46	19.7	13.6-25.8	14	8.1	03.4-12.8	4	2.0	00.0-04.0	38	19.0	12.9-25.1	2	0.9	00.0-02.1
Non-Native	199	18,514	101	48.9	40.7-57.1	43	19.9	13.4-26.4	14	8.8	03.7-13.9	4	2.2	00.0-04.4	35	19.4	12.9-25.9	2	0.9	00.0-02.3
55-64	68	5,810	39	57.8	44.5-71.1	14	20.2	08.8-31.6	7	10.7	02.5-18.9	2	4.3	00.0-10.2	6	7.1	01.2-13.0			
65+	113	8,994	56	47.6	37.4-57.8	24	18.3	11.0-25.6	4	4.2	00.3-08.1	2	1.7	00.0-04.1	25	26.4	16.8-36.0	2	1.9	00.0-04.4
H.S. or G.E.D.	66	5,843	33	46.3	32.6-60.0	16	25.7	13.2-38.2	2	3.9	00.0-09.4	4	6.8	00.1-13.5	9	14.4	04.8-24.0	2	2.9	00.0-07.0
Some Post-H.S.	63	6,868	36	52.3	38.2-66.4	8	14.4	04.4-24.4	10	17.5	05.7-29.3				9	15.8	05.6-26.0			
College Graduate	57	3,810	27	48.6	33.1-64.1	20	33.2	19.3-47.1	2	5.2	00.0-13.0				8	13.0	03.6-22.4			

NOTE:

^{&#}x27;% = Percentage, C.I.(95%) = Confidence Interval (at 95 percent probability level). Percentages are weighted to population characteristics.

Rows with less than 50 observations have been suppressed.

TABLE 12: ABOUT HOW OFTEN DO YOU CHECK YOUR FEET FOR ANY SORES OR IRRITATIONS? INCLUDE TIMES WHEN CHECKED BY A FAMILY MEMBER OR FRIEND, BUT DO NOT INCLUDE TIMES WHEN CHECKED BY A HEALTH PROFESSIONAL. (FEETCHK2)

Denominator excludes: Respondents who are male or respondents with do not know/refused or with missing responses Denominator excludes: Respondents that reported not being diagnosed with diabetes Denominator excludes: Women respondents that reported being diagnosed with diabetes only during pregnancy

Data was weighted using _LANDWT.

DEMOGRAPHIC	RESPONDE	NT NUMBER	Т	imes p	er Day	Tir	mes pe	r Week	Tin	nes pe	r Month	Ti	mes p	er Year		Nev	er		No F	eet
GROUPS	TOTAL	WEIGHTED	N	%	C.I. (95%)	N	%	C.I. (95%)	N	%	C.I. (95%)	N	%	C.I. (95%)	N	%	C.I. (95%)	N	%	C.I. (95%)
Female	279	19,978	170	62.9	55.8-70.0	52	17.8	12.3-23.3	19	6.9	02.8-11.0	1	0.2	00.0-00.6	35	11.5	07.2-15.8	2	0.7	00.0-01.7
Non-Native	259	17,324	154	61.1	54.0-68.2	48	16.7	11.8-21.6	19	8.0	03.3-12.7	1	0.2	00.0-00.6	35	13.2	08.5-17.9	2	0.8	00.0-02.0
55-64	70	5,345	53	79.8	69.6-90.0	10	11.3	03.5-19.1	3	4.1	00.0-08.8	1	0.8	00.0-02.4	3	4.1	00.0-09.0			
65+	161	9,026	88	54.8	46.2-63.4	34	22.0	14.9-29.1	11	6.2	02.3-10.1				26	15.5	09.2-21.8	2	1.6	00.0-03.8
H.S. or G.E.D.	112	7,306	70	62.7	51.3-74.1	19	19.4	08.8-30.0	9	6.0	01.7-10.3	1	0.6	00.0-01.8	13	11.3	04.2-18.4			
Some Post-H.S.	92	8,135	55	64.2	52.8-75.6	18	15.0	07.7-22.3	6	9.0	00.2-17.8				11	10.1	03.8-16.4	2	1.7	00.0-04.1
College Graduate	50	2,420	31	60.0	43.5-76.5	10	23.4	07.9-38.9	3	6.9	00.0-14.9				6	9.7	01.9-17.5			
\$15,000- 24,999	63	4,378	37	58.3	42.2-74.4	14	20.4	08.8-32.0	4	11.3	00.0-25.8				7	8.3	01.6-15.0	1	1.6	00.0-04.7

NOTE:

^{&#}x27;% = Percentage, C.I.(95%) = Confidence Interval (at 95 percent probability level). Percentages are weighted to population characteristics. Rows with less than 50 observations have been suppressed.

MODULE 2: DIABETES

TABLE 13: ABOUT HOW MANY TIMES IN THE PAST 12 MONTHS HAVE YOU SEEN A DOCTOR, NURSE, OR OTHER HEALTH PROFESSIONAL FOR YOUR DIABETES? (DOCTDIAB)

Denominator excludes: Respondents with do not know/refused/missing responses

Denominator excludes: Respondents that reported not being diagnosed with diabetes

Denominator excludes: Women respondents that reported being diagnosed with diabetes only during pregnancy

Data was weighted using LANDWT

DEMOGRAPHIC	RESPONDE	NT NUMBER		1 tim	ne		2 tim	es		3 tim	es		4 tim	es	5 0	or more	times		Non	е
GROUPS	TOTAL	WEIGHTED	N	%	C.I. (95%)	N	%	C.I. (95%)	N	%	C.I. (95%)	N	%	C.I. (95%)	N	%	C.I. (95%)	N	%	C.I. (95%)
TOTAL	491	40,401	95	17.5	13.8-21.2	133	27.9	23.0-32.8	75	15.3	11.6-19.0	92	19.1	14.8-23.4	43	10.9	07.2-14.6	53	9.3	06.4-12.2
Male	218	20,813	40	17.2	11.5-22.9	70	34.7	27.1-42.3	29	14.4	08.9-19.9	28	12.0	07.3-16.7	24	11.4	06.5-16.3	27	10.3	06.2-14.4
Female	273	19,588	55	17.8	12.9-22.7	63	20.7	15.4-26.0	46	16.2	11.3-21.1	64	26.6	19.7-33.5	19	10.4	05.1-15.7	26	8.3	04.2-12.4
Non-Native	455	36,159	92	19.1	15.0-23.2	129	30.2	25.1-35.3	70	16.1	12.2-20.0	82	16.7	12.8-20.6	31	8.0	04.7-11.3	51	9.9	06.8-13.0
45-54	63	7,678	14	13.7	06.1-21.3	16	34.9	20.4-49.4	15	23.3	11.7-34.9	12	17.3	07.3-27.3	5	9.7	01.3-18.1	1	1.2	00.0-03.6
55-64	137	11,136	23	16.0	08.9-23.1	44	28.7	20.5-36.9	19	13.0	06.7-19.3	22	18.4	09.4-27.4	17	14.7	06.9-22.5	12	9.3	03.8-14.8
65+	272	18,165	55	20.9	15.4-26.4	68	25.2	19.5-30.9	40	15.3	10.4-20.2	54	19.2	14.1-24.3	18	8.1	03.8-12.4	37	11.3	07.4-15.2
Less Than H.S.	55	6,041	11	19.9	07.9-31.9	14	21.7	10.3-33.1	9	16.8	05.6-28.0	11	21.8	07.5-36.1	7	15.6	03.3-27.9	3	4.2	00.0-09.1
H.S. or G.E.D.	173	12,949	37	19.2	12.7-25.7	43	25.2	17.4-33.0	30	17.8	10.7-24.9	33	21.1	13.3-28.9	11	8.0	02.7-13.3	19	8.7	04.4-13.0
Some Post-H.S.	156	15,192	30	16.6	10.3-22.9	45	30.7	22.1-39.3	22	12.5	07.0-18.0	28	17.6	10.9-24.3	14	11.3	05.0-17.6	17	11.3	05.2-17.4
College Graduate	106	6,178	16	13.2	06.5-19.9	31	32.8	21.4-44.2	14	15.6	07.0-24.2	20	16.0	08.4-23.6	11	11.4	03.8-19.0	14	10.9	05.0-16.8
Less than \$15,000	61	5,814	10	11.0	03.7-18.3	12	15.2	05.2-25.2	11	18.2	06.6-29.8	16	34.3	18.0-50.6	7	15.3	04.1-26.5	5	6.0	00.7-11.3
\$15,000- 24,999	90	7,177	18	19.8	09.2-30.4	23	23.3	13.7-32.9	12	9.8	03.9-15.7	20	21.4	11.0-31.8	8	15.6	04.2-27.0	9	10.0	00.4-19.6
\$25,000- 34,999	75	5,894	10	13.0	04.4-21.6	28	39.4	26.1-52.7	11	15.1	05.7-24.5	10	12.6	03.6-21.6	7	5.8	01.3-10.3	9	14.1	04.5-23.7
\$35,000- 49,999	64	5,321	12	14.7	06.1-23.3	19	38.6	23.9-53.3	5	7.3	00.6-14.0	17	23.0	11.6-34.4	6	11.7	01.5-21.9	5	4.6	00.3-08.9
\$50,000- 74,999	58	5,424	13	22.0	10.2-33.8	20	40.3	24.6-56.0	11	16.7	06.5-26.9	7	8.7	01.6-15.8	3	6.8	00.0-15.6	4	5.4	00.1-10.7
\$75,000+	57	4,546	14	21.8	10.0-33.6	10	15.5	04.7-26.3	12	28.9	13.4-44.4	8	12.3	03.5-21.1	3	6.2	00.0-13.6	10	15.2	05.6-24.8

NOTE:

^{&#}x27;% = Percentage, C.I.(95%) = Confidence Interval (at 95 percent probability level). Percentages are weighted to population characteristics. Rows with less than 50 observations have been suppressed.

TABLE 14: ABOUT HOW MANY TIMES IN THE PAST 12 MONTHS HAVE YOU SEEN A DOCTOR, NURSE, OR OTHER HEALTH PROFESSIONAL FOR YOUR DIABETES? (DOCTDIAB)

Denominator excludes: Respondents who are female or respondents with do not know/refused or with missing responses

Denominator excludes: Respondents that reported not being diagnosed with diabetes

Denominator excludes: Women respondents that reported being diagnosed with diabetes only during pregnancy

Data was weighted using _LANDWT.

DEMOGRAPHIC	RESPONDE	NT NUMBER		1 tin	ne		2 tim	ies		3 tim	es		4 tim	es	5 (or more	times		Non	е
GROUPS	TOTAL	WEIGHTED	N	%	C.I. (95%)	N	%	C.I. (95%)	N	%	C.I. (95%)									
Male	218	20,813	40	17.2	11.5-22.9	70	34.7	27.1-42.3	29	14.4	08.9-19.9	28	12.0	07.3-16.7	24	11.4	06.5-16.3	27	10.3	06.2-14.4
Non-Native	202	19,225	39	18.4	12.3-24.5	68	36.8	28.8-44.8	25	14.0	08.1-19.9	26	11.2	06.7-15.7	19	9.4	04.5-14.3	25	10.2	05.9-14.5
55-64	69	5,999	10	14.7	04.5-24.9	24	32.2	20.4-44.0	8	10.2	02.9-17.5	6	9.6	01.2-18.0	11	18.3	06.7-29.9	10	15.0	05.8-24.2
65+	114	9,286	27	25.3	16.1-34.5	30	23.9	15.7-32.1	13	14.1	06.3-21.9	19	16.3	09.0-23.6	10	9.7	03.2-16.2	15	10.7	05.0-16.4
H.S. or G.E.D.	66	5,960	15	18.4	09.2-27.6	23	37.8	24.1-51.5	6	10.6	00.6-20.6	8	12.5	02.7-22.3	7	11.1	02.9-19.3	7	9.5	02.2-16.8
Some Post-H.S.	64	7,097	11	16.4	06.2-26.6	24	42.9	28.6-57.2	10	13.6	05.0-22.2	6	8.8	01.9-15.7	5	6.3	00.2-12.4	8	12.0	03.4-20.6
College Graduate	56	3,759	9	14.1	04.7-23.5	14	28.2	12.7-43.7	8	18.4	05.9-30.9	9	15.5	04.7-26.3	7	11.0	02.4-19.6	9	12.8	04.2-21.4

NOTE:

^{&#}x27;% = Percentage, C.I.(95%) = Confidence Interval (at 95 percent probability level). Percentages are weighted to population characteristics.

Rows with less than 50 observations have been suppressed.

MODULE 2: DIABETES

TABLE 15: ABOUT HOW MANY TIMES IN THE PAST 12 MONTHS HAVE YOU SEEN A DOCTOR, NURSE, OR OTHER HEALTH PROFESSIONAL FOR YOUR DIABETES? (DOCTDIAB)

Denominator excludes: Respondents who are male or respondents with do not know/refused or with missing responses

Denominator excludes: Respondents that reported not being diagnosed with diabetes

Denominator excludes: Women respondents that reported being diagnosed with diabetes only during pregnancy

Data was weighted using _LANDWT.

DEMOGRAPHIC	RESPONDE	NT NUMBER		1 tin	ne		2 tim	es		3 tim	es		4 tim	ies	5 0	or more	e times		Non	ne
GROUPS	TOTAL	WEIGHTED	N	%	C.I. (95%)	N	%	C.I. (95%)	N	%	C.I. (95%)									
Female	273	19,588	55	17.8	12.9-22.7	63	20.7	15.4-26.0	46	16.2	11.3-21.1	64	26.6	19.7-33.5	19	10.4	05.1-15.7	26	8.3	04.2-12.4
Non-Native	253	16,934	53	19.8	14.3-25.3	61	22.8	16.9-28.7	45	18.4	12.9-23.9	56	23.0	16.9-29.1	12	6.5	02.0-11.0	26	9.6	04.9-14.3
55-64	68	5,137	13	17.4	07.4-27.4	20	24.5	13.3-35.7	11	16.3	06.3-26.3	16	28.7	13.0-44.4	6	10.5	00.7-20.3	2	2.6	00.0-06.3
65+	158	8,878	28	16.2	10.3-22.1	38	26.6	18.8-34.4	27	16.5	10.2-22.8	35	22.3	15.2-29.4	8	6.5	01.0-12.0	22	11.9	06.8-17.0
H.S. or G.E.D.	107	6,989	22	19.8	11.0-28.6	20	14.4	07.7-21.1	24	24.0	14.4-33.6	25	28.3	17.1-39.5	4	5.4	00.0-12.1	12	8.0	03.1-12.9
Some Post-H.S.	92	8,095	19	16.8	09.0-24.6	21	20.0	11.4-28.6	12	11.5	04.6-18.4	22	25.2	14.6-35.8	9	15.7	05.5-25.9	9	10.7	01.9-19.5
College Graduate	50	2,420	7	11.8	02.8-20.8	17	40.0	23.5-56.5	6	11.3	02.1-20.5	11	16.8	06.8-26.8	4	12.1	00.0-26.4	5	7.9	00.5-15.3
\$15,000- 24,999	60	4.170	11	13.8	04.8-22.8	13	19.9	08.1-31.7	9	11.8	03.8-19.8	16	29.5	14.4-44.6	3	10.7	00.0-23.4	8	14.4	00.0-29.3

^{&#}x27;% = Percentage, C.I.(95%) = Confidence Interval (at 95 percent probability level). Percentages are weighted to population characteristics. Rows with less than 50 observations have been suppressed.

MODULE 2: DIABETES

TABLE 16: A TEST FOR "A ONE C" MEASURES THE AVERAGE LEVEL OF BLOOD SUGAR OVER THE PAST THREE MONTHS. ABOUT HOW MANY TIMES IN THE PAST 12 MONTHS HAS A DOCTOR, NURSE, OR OTHER HEALTH PROFESSIONAL CHECKED YOU FOR "A ONE C"? (CHKHEMO3)

> Denominator excludes: Respondents with do not know/refused/missing responses Denominator excludes: Respondents that reported not being diagnosed with diabetes Denominator excludes: Women respondents that reported being diagnosed with diabetes only during pregnancy

> > Data was weighted using _LANDWT.

										_										
DEMOGRAPHIC	RESPONDE	NT NUMBER		1-2 pe	r yr		3-4 pe	er yr		5-9 pe	er yr	10 (or mor	e per yr		NON	ΙE	Ne	ever He	eard of
GROUPS	TOTAL	WEIGHTED	N	%	C.I. (95%)	N	%	C.I. (95%)	N	%	C.I. (95%)	N	%	C.I. (95%)	N	%	C.I. (95%)	N	%	C.I. (95%)
TOTAL	473	38,542	251	52.1	46.6-57.6	154	33.6	28.5-38.7	11	3.0	00.8-05.2	4	0.6	00.0-01.2	28	6.2	03.5-08.9	25	4.5	02.5-06.5
Male	204	19,119	120	59.4	51.8-67.0	55	25.2	18.5-31.9	4	3.3	00.0-06.6	2	0.7	00.0-01.9	17	8.6	04.3-12.9	6	2.7	00.3-05.1
Female	269	19,422	131	44.9	37.8-52.0	99	41.8	34.4-49.2	7	2.7	00.2-05.2	2	0.5	00.0-01.3	11	3.9	00.4-07.4	19	6.1	03.2-09.0
Non-Native	439	34,463	241	56.0	50.5-61.5	136	29.9	24.8-35.0	8	2.0	00.2-03.8	4	0.7	00.0-01.5	28	7.0	03.9-10.1	22	4.4	02.4-06.4
45-54	60	7,130	35	56.3	42.0-70.6	18	32.5	19.0-46.0	4	7.7	00.0-15.7				1	2.1	00.0-06.2	2	1.3	00.0-03.3
55-64	137	11,170	77	54.3	44.3-64.3	49	35.3	25.5-45.1	2	2.1	00.0-05.2				6	6.1	01.2-11.0	3	2.2	00.0-04.7
65+	257	16,820	133	53.3	46.4-60.2	80	30.9	24.6-37.2	5	2.2	00.0-04.9	4	1.5	00.0-03.1	19	6.2	03.3-09.1	16	5.9	02.8-09.0
Less Than H.S.	51	5,484	28	52.7	36.6-68.8	14	31.5	15.6-47.4	1	3.9	00.0-11.3				2	3.5	00.0-08.4	6	8.4	01.1-15.7
H.S. or G.E.D.	164	12,354	84	50.1	40.9-59.3	52	33.3	24.3-42.3	5	4.9	00.2-09.6	2	0.6	00.0-01.4	14	6.3	02.4-10.2	7	4.7	01.0-08.4
Some Post-H.S.	152	14,573	83	52.2	43.0-61.4	48	33.1	24.5-41.7	2	1.6	00.0-04.0	2	1.2	00.0-02.8	10	9.0	02.9-15.1	7	2.9	00.5-05.3
College Graduate	104	6,056	55	55.4	44.0-66.8	39	37.1	26.1-48.1	3	1.5	00.0-03.3				2	1.8	00.0-04.3	5	4.2	00.5-07.9
Less than \$15,000	57	5,541	26	36.8	22.3-51.3	17	41.0	24.3-57.7	4	8.8	00.0-18.2	1	1.3	00.0-03.8	5	7.0	00.7-13.3	4	5.1	00.0-10.2
\$15,000- 24,999	84	6,618	44	50.5	37.0-64.0	30	36.7	23.8-49.6	2	2.9	00.0-07.8				6	8.6	00.0-18.6	2	1.4	00.0-03.4
\$25,000- 34,999	74	5,857	43	59.0	45.7-72.3	20	26.2	14.4-38.0	2	1.2	00.0-03.0	3	2.9	00.0-06.6	4	8.1	00.0-16.5	2	2.5	00.0-06.0
\$35,000- 49,999	63	4,987	31	51.0	36.7-65.3	25	36.7	23.4-50.0	1	4.6	00.0-13.4				5	6.8	00.1-13.5	1	0.8	00.0-02.4
\$50,000- 74,999	56	5,072	36	67.8	53.7-81.9	17	28.3	14.6-42.0	1	1.4	00.0-04.1							2	2.5	00.0-06.0
\$75,000+	56	4,465	31	52.2	36.7-67.7	20	37.1	21.8-52.4							2	4.6	00.0-11.3	3	6.1	00.0-13.9

NOTE:

^{&#}x27;% = Percentage, C.I.(95%) = Confidence Interval (at 95 percent probability level). Percentages are weighted to population characteristics. Rows with less than 50 observations have been suppressed.

MODULE 2: DIABETES

TABLE 17: A TEST FOR "A ONE C" MEASURES THE AVERAGE LEVEL OF BLOOD SUGAR OVER THE PAST THREE MONTHS. ABOUT HOW MANY TIMES IN THE PAST 12 MONTHS HAS A DOCTOR, NURSE, OR OTHER HEALTH PROFESSIONAL CHECKED YOU FOR "A ONE C"? (CHKHEMO3)

Denominator excludes: Respondents who are female or respondents with do not know/refused or with missing responses

Denominator excludes: Respondents that reported not being diagnosed with diabetes

Denominator excludes: Women respondents that reported being diagnosed with diabetes only during pregnancy

Data was weighted using LANDWT.

DEMOGRAPHIC	RESPONDE	NT NUMBER		1-2 pe	er yr		3-4 pe	r yr		5-9 pe		10 c	or mor	e per yr		NON	ΙE	Ne	ever He	ard of
GROUPS	TOTAL	WEIGHTED	N	%	C.I. (95%)	N	%	C.I. (95%)	N	%	C.I. (95%)	N	%	C.I. (95%)	N	%	C.I. (95%)	N	%	C.I. (95%)
Male	204	19,119	120	59.4	51.8-67.0	55	25.2	18.5-31.9	4	3.3	00.0-06.6	2	0.7	00.0-01.9	17	8.6	04.3-12.9	6	2.7	00.3-05.1
Non-Native	189	17,596	115	62.3	54.5-70.1	47	22.3	15.8-28.8	3	2.7	00.0-05.8	2	0.8	00.0-02.0	17	9.3	04.6-14.0	5	2.6	00.2-05.0
55-64	67	5,824	41	61.7	48.6-74.8	19	23.8	13.2-34.4	1	2.8	00.0-08.3				6	11.7	02.5-20.9			
65+	105	8,316	61	55.8	45.4-66.2	28	27.7	18.3-37.1	1	2.6	00.0-07.7	2	1.7	00.0-04.2	9	7.6	02.5-12.7	4	4.6	00.1-09.1
H.S. or G.E.D.	61	5,587	40	63.0	48.7-77.3	11	20.5	07.8-33.2	2	4.6	00.0-10.9	1	0.8	00.0-02.4	5	8.0	00.4-15.6	2	3.1	00.0-07.4
Some Post-H.S.	60	6,430	33	58.4	44.7-72.1	15	21.4	10.8-32.0	1	2.6	00.0-07.5	1	1.5	00.0-04.4	8	14.0	04.4-23.6	2	2.1	00.0-05.0
College Graduate	55	3,678	30	56.4	41.1-71.7	22	38.6	23.7-53.5							2	2.9	00.0-07.0	1	2.0	00.0-05.9

NOTE:

^{&#}x27;% = Percentage, C.I.(95%) = Confidence Interval (at 95 percent probability level). Percentages are weighted to population characteristics.

Rows with less than 50 observations have been suppressed.

MODULE 2: DIABETES

TABLE 18: A TEST FOR "A ONE C" MEASURES THE AVERAGE LEVEL OF BLOOD SUGAR OVER THE PAST THREE MONTHS. ABOUT HOW MANY TIMES IN THE PAST 12 MONTHS HAS A DOCTOR, NURSE, OR OTHER HEALTH PROFESSIONAL CHECKED YOU FOR "A ONE C"? (CHKHEMO3)

Denominator excludes: Respondents who are male or respondents with do not know/refused or with missing responses

Denominator excludes: Respondents that reported not being diagnosed with diabetes

Denominator excludes: Women respondents that reported being diagnosed with diabetes only during pregnancy

Data was weighted using _LANDWT.

DEMOGRAPHIC	RESPONDE	NT NUMBER		1-2 pe	er yr		3-4 pe	r yr		5-9 pe	er yr	10	or mor	e per yr		NON	1E	Ne	ever He	eard of
GROUPS	TOTAL	WEIGHTED	N	%	C.I. (95%)	N	%	C.I. (95%)	N	%	C.I. (95%)	N	%	C.I. (95%)	N	%	C.I. (95%)	N	%	C.I. (95%)
Female	269	19,422	131	44.9	37.8-52.0	99	41.8	34.4-49.2	7	2.7	00.2-05.2	2	0.5	00.0-01.3	11	3.9	00.4-07.4	19	6.1	03.2-09.0
Non-Native	250	16,868	126	49.4	42.1-56.7	89	37.9	30.6-45.2	5	1.3	00.1-02.5	2	0.6	00.0-01.6	11	4.5	00.4-08.6	17	6.3	03.0-09.6
55-64	70	5,345	36	46.2	31.9-60.5	30	47.9	33.0-62.8	1	1.3	00.0-03.8							3	4.6	00.0-09.7
65+	152	8,504	72	50.8	42.0-59.6	52	34.1	25.7-42.5	4	1.8	00.0-03.8	2	1.2	00.0-03.0	10	4.9	01.6-08.2	12	7.2	03.1-11.3
H.S. or G.E.D.	103	6,768	44	39.4	28.2-50.6	41	43.9	31.7-56.1	3	5.2	00.0-12.1	1	0.5	00.0-01.5	9	4.9	01.4-08.4	5	6.0	00.1-11.9
Some Post-H.S.	92	8,143	50	47.3	35.5-59.1	33	42.2	30.0-54.4	1	0.9	00.0-02.7	1	0.9	00.0-02.7	2	5.2	00.0-13.2	5	3.6	00.3-06.9
\$15,000- 24,999	58	4,062	25	36.1	21.2-51.0	26	50.3	33.4-67.2	1	0.6	00.0-01.8				4	10.8	00.0-26.1	2	2.3	00.0-05.4

NOTE:

^{&#}x27;% = Percentage, C.I.(95%) = Confidence Interval (at 95 percent probability level). Percentages are weighted to population characteristics.

Rows with less than 50 observations have been suppressed.

MODULE 2: DIABETES

TABLE 19: ABOUT HOW MANY TIMES IN THE PAST 12 MONTHS HAS A HEALTH PROFESSIONAL CHECKED YOUR FEET FOR ANY SORES OR IRRITATIONS? (FEETCHK)

Denominator excludes: Respondents with do not know/refused/missing responses

Denominator excludes: Respondents that reported not being diagnosed with diabetes

Denominator excludes: Respondents that reported having no feet

Denominator excludes: Women respondents that reported being diagnosed with diabetes only during pregnancy

Data was weighted using _LANDWT.

DEMOGRAPHIC	RESPONDE	NT NUMBER		1 tim	ne		2 tim	es		3 tim	es		4 tim	es	5 o	r more	times		Non	e
GROUPS	TOTAL	WEIGHTED	N	%	C.I. (95%)	N	%	C.I. (95%)	N	%	C.I. (95%)	N	%	C.I. (95%)	N	%	C.I. (95%)	N	%	C.I. (95%)
TOTAL	490	40,068	140	27.3	22.6-32.0	111	24.2	19.5-28.9	53	12.5	08.6-16.4	64	12.7	09.4-16.0	28	4.6	02.6-06.6	94	18.7	14.6-22.8
Male	213	20,188	62	28.5	21.4-35.6	61	29.3	22.0-36.6	19	10.3	05.2-15.4	19	8.8	04.5-13.1	14	5.2	02.5-07.9	38	17.8	11.7-23.9
Female	277	19,880	78	26.0	19.9-32.1	50	19.1	13.0-25.2	34	14.8	08.9-20.7	45	16.6	11.3-21.9	14	4.0	01.5-06.5	56	19.5	13.8-25.2
Non-Native	456	36,013	135	28.8	23.9-33.7	104	24.5	19.8-29.2	49	11.7	08.0-15.4	56	11.5	08.2-14.8	21	3.3	01.7-04.9	91	20.3	15.8-24.8
45-54	63	7,678	16	20.8	09.4-32.2	18	30.8	17.7-43.9	8	15.4	04.6-26.2	7	11.8	03.0-20.6	2	2.4	00.0-05.7	12	18.8	07.8-29.8
55-64	139	11,256	38	25.1	16.7-33.5	34	25.4	16.4-34.4	19	14.5	07.1-21.9	20	14.6	07.5-21.7	7	4.3	01.0-07.6	21	16.0	09.1-22.9
65+	270	17,895	82	33.1	26.6-39.6	57	22.5	16.8-28.2	24	8.8	05.1-12.5	35	12.1	08.0-16.2	18	5.5	02.8-08.2	54	18.0	12.9-23.1
Less Than H.S.	55	6,020	13	25.0	12.1-37.9	16	32.1	16.8-47.4	8	14.9	03.5-26.3	4	6.0	00.1-11.9	3	3.9	00.0-08.4	11	18.0	06.6-29.4
H.S. or G.E.D.	173	12,922	48	25.7	18.3-33.1	36	23.4	15.6-31.2	21	15.4	07.8-23.0	22	13.2	07.1-19.3	11	4.8	01.7-07.9	35	17.4	11.1-23.7
Some Post-H.S.	153	14,823	50	31.7	23.1-40.3	33	22.1	14.7-29.5	14	9.1	03.4-14.8	22	15.4	08.7-22.1	4	2.5	00.0-05.0	30	19.2	11.9-26.5
College Graduate	107	6,230	28	21.9	13.7-30.1	26	23.7	14.5-32.9	10	12.4	04.0-20.8	15	11.4	05.5-17.3	10	9.9	02.8-17.0	18	20.8	10.2-31.4
Less than \$15,000	61	5,781	16	22.8	10.6-35.0	12	27.6	11.5-43.7	3	10.0	00.0-22.5	10	14.3	04.9-23.7	6	7.4	01.1-13.7	14	17.8	08.0-27.6
\$15,000- 24,999	90	7,212	24	24.8	14.0-35.6	14	13.3	06.0-20.6	19	23.5	11.9-35.1	12	13.1	04.3-21.9	7	6.8	00.5-13.1	14	18.6	07.2-30.0
\$25,000- 34,999	73	5,633	22	26.6	15.2-38.0	19	25.7	13.9-37.5	8	11.0	02.6-19.4	6	10.2	01.2-19.2	5	5.0	00.1-09.9	13	21.4	08.9-33.9
\$35,000- 49,999	65	5,354	17	30.0	16.5-43.5	17	28.5	16.0-41.0	7	16.9	03.8-30.0	8	9.4	02.7-16.1	5	5.1	00.4-09.8	11	10.0	03.7-16.3
\$50,000- 74,999	58	5,329	17	30.4	15.7-45.1	18	29.8	15.7-43.9	5	7.4	00.7-14.1	11	17.8	06.4-29.2				7	14.5	03.7-25.3
\$75,000+	57	4,546	18	26.7	14.5-38.9	14	30.0	14.9-45.1	6	10.3	00.9-19.7	6	9.4	01.4-17.4				13	23.6	10.5-36.7

NOTE:

^{&#}x27;% = Percentage, C.I.(95%) = Confidence Interval (at 95 percent probability level). Percentages are weighted to population characteristics. Rows with less than 50 observations have been suppressed.

MODULE 2: DIABETES

TABLE 20: ABOUT HOW MANY TIMES IN THE PAST 12 MONTHS HAS A HEALTH PROFESSIONAL CHECKED YOUR FEET FOR ANY SORES OR IRRITATIONS? (FEETCHK)

Denominator excludes: Respondents who are female or respondents with do not know/refused or with missing responses

Denominator excludes: Respondents that reported not being diagnosed with diabetes

Denominator excludes: Respondents that reported having no feet

Denominator excludes: Women respondents that reported being diagnosed with diabetes only during pregnancy

Data was weighted using _LANDWT.

DEMOGRAPHIC	RESPONDE	NT NUMBER		1 tin	ne		2 tim	es		3 tim	es		4 tim	es	5 c	r more	times		Non	е
GROUPS	TOTAL	WEIGHTED	N	%	C.I. (95%)	N	%	C.I. (95%)	N	%	C.I. (95%)									
Male	213	20,188	62	28.5	21.4-35.6	61	29.3	22.0-36.6	19	10.3	05.2-15.4	19	8.8	04.5-13.1	14	5.2	02.5-07.9	38	17.8	11.7-23.9
Non-Native	199	18,787	60	29.7	22.3-37.1	59	30.6	23.0-38.2	17	9.9	04.6-15.2	17	7.4	03.7-11.1	10	3.8	01.3-06.3	36	18.5	12.0-25.0
55-64	68	5,877	17	24.8	12.8-36.8	20	27.4	16.0-38.8	9	14.1	03.1-25.1	5	9.0	00.4-17.6	5	5.7	00.6-10.8	12	19.0	08.6-29.4
65+	111	8,967	39	37.1	27.1-47.1	29	24.2	15.6-32.8	5	5.7	00.8-10.6	12	10.0	04.3-15.7	7	6.0	01.5-10.5	19	17.0	09.0-25.0
H.S. or G.E.D.	63	5,724	19	26.6	14.4-38.8	24	40.4	26.3-54.5	2	3.2	00.0-07.7	5	10.6	00.6-20.6	4	6.2	00.1-12.3	9	12.9	04.3-21.5
Some Post-H.S.	62	6,779	22	33.9	20.4-47.4	16	27.5	14.6-40.4	5	9.0	00.2-17.8	6	9.7	01.9-17.5	2	2.8	00.0-06.5	11	17.1	06.9-27.3
College Graduate	57	3,810	12	18.5	07.9-29.1	13	21.6	09.6-33.6	8	17.6	04.9-30.3	7	10.4	02.6-18.2	6	8.6	01.5-15.7	11	23.3	08.4-38.2

NOTE:

^{&#}x27;% = Percentage, C.I.(95%) = Confidence Interval (at 95 percent probability level). Percentages are weighted to population characteristics. Rows with less than 50 observations have been suppressed.

MODULE 2: DIABETES

TABLE 21: ABOUT HOW MANY TIMES IN THE PAST 12 MONTHS HAS A HEALTH PROFESSIONAL CHECKED YOUR FEET FOR ANY SORES OR IRRITATIONS? (FEETCHK)

Denominator excludes: Respondents who are male or respondents with do not know/refused or with missing responses

Denominator excludes: Respondents that reported not being diagnosed with diabetes

Denominator excludes: Respondents that reported having no feet

Denominator excludes: Women respondents that reported being diagnosed with diabetes only during pregnancy

Data was weighted using _LANDWT.

DEMOGRAPHIC	RESPONDE	NT NUMBER		1 tin	ne		2 tim	es		3 tim	es		4 tim	es	5 c	r more	times		Non	e
GROUPS	TOTAL	WEIGHTED	N	%	C.I. (95%)	N	%	C.I. (95%)	N	%	C.I. (95%)									
Female	277	19,880	78	26.0	19.9-32.1	50	19.1	13.0-25.2	34	14.8	08.9-20.7	45	16.6	11.3-21.9	14	4.0	01.5-06.5	56	19.5	13.8-25.2
Non-Native	257	17,226	75	27.8	21.5-34.1	45	17.8	12.5-23.1	32	13.6	08.5-18.7	39	16.0	10.7-21.3	11	2.7	00.9-04.5	55	22.1	15.8-28.4
55-64	71	5,379	21	25.5	13.7-37.3	14	23.2	08.5-37.9	10	15.0	05.2-24.8	15	20.8	09.6-32.0	2	2.8	00.0-07.1	9	12.7	03.9-21.5
65+	159	8,928	43	29.0	20.8-37.2	28	20.8	13.4-28.2	19	11.9	06.4-17.4	23	14.1	08.2-20.0	11	5.1	02.0-08.2	35	19.1	13.0-25.2
H.S. or G.E.D.	110	7,198	29	25.0	15.8-34.2	12	9.9	03.8-16.0	19	25.1	13.1-37.1	17	15.2	07.8-22.6	7	3.8	00.9-06.7	26	21.0	12.2-29.8
Some Post-H.S.	91	8,044	28	29.9	19.1-40.7	17	17.5	09.1-25.9	9	9.2	01.9-16.5	16	20.2	10.0-30.4	2	2.3	00.0-05.4	19	21.0	10.8-31.2
College Graduate	50	2,420	16	27.2	14.1-40.3	13	26.9	12.6-41.2	2	4.2	00.0-10.1	8	13.0	04.0-22.0	4	11.9	00.0-26.0	7	16.8	03.1-30.5
\$15.000- 24.999	61	4.270	14	20.3	08.5-32.1	7	8.7	01.6-15.8	15	26.1	11.8-40.4	9	16.2	03.9-28.5	5	7.1	00.0-15.7	11	21.5	06.0-37.0

NOTE:

^{&#}x27;% = Percentage, C.I.(95%) = Confidence Interval (at 95 percent probability level). Percentages are weighted to population characteristics. Rows with less than 50 observations have been suppressed.

MODULE 2: DIABETES

TABLE 22: WHEN WAS THE LAST TIME YOU HAD AN EYE EXAM IN WHICH THE PUPILS WERE DILATED? THIS WOULD HAVE MADE YOU TEMPORARILY SENSITIVE TO BRIGHT LIGHT. (EYEEXAM)

Denominator excludes: Respondents with do not know/refused/missing responses

Denominator excludes: Respondents that reported not being diagnosed with diabetes

Denominator excludes: Women respondents that reported being diagnosed with diabetes only during pregnancy

Data was weighted using _LANDWT.

DEMOCD ADUID	DECDONDE	NT NUMBER		4 14 4	L A	4 14	J	40 Mandha	4 . V		0 V	•		V		NI	
DEMOGRAPHIC		NT NUMBER	<	1 Mont	-	1 Wont		12 Months	1+ Y	rs but •	< 2 Years		or more			Nev	
GROUPS	TOTAL	WEIGHTED	N	%	C.I. (95%)	N	%	C.I. (95%)	N	%	C.I. (95%)	N	%	C.I. (95%)	N	%	C.I. (95%)
TOTAL	498	40,503	79	14.6	10.9-18.3	277	54.0	48.7-59.3	74	16.7	12.4-21.0	58	12.1	08.6-15.6	10	2.6	00.8-04.4
Male	217	20,552	30	12.4	07.7-17.1	120	52.8	45.0-60.6	30	16.0	09.7-22.3	29	14.1	08.4-19.8	8	4.6	01.1-08.1
Female	281	19,951	49	16.9	11.2-22.6	157	55.3	48.0-62.6	44	17.4	11.5-23.3	29	10.0	05.9-14.1	2	0.5	00.0-01.1
Non-Native	463	36,331	71	13.3	09.8-16.8	254	52.7	47.2-58.2	72	18.0	13.3-22.7	56	13.1	09.2-17.0	10	2.9	00.9-04.9
45-54	62	7,603	5	8.3	00.9-15.7	32	45.5	31.4-59.6	13	25.2	11.7-38.7	10	17.5	06.7-28.3	2	3.6	00.0-08.5
55-64	138	11,188	20	16.0	08.4-23.6	76	55.2	45.2-65.2	21	13.8	07.7-19.9	18	13.0	06.1-19.9	3	2.0	00.0-04.4
65+	278	18,257	50	15.7	11.2-20.2	162	60.7	54.2-67.2	36	13.6	08.7-18.5	26	8.4	05.1-11.7	4	1.5	00.0-03.1
Less Than H.S.	56	6,016	4	4.4	00.0-08.9	38	67.1	52.6-81.6	6	12.5	02.1-22.9	6	12.0	01.2-22.8	2	4.0	00.0-09.5
H.S. or G.E.D.	178	13,117	32	17.2	10.1-24.3	84	44.5	35.9-53.1	30	15.7	09.8-21.6	28	19.1	11.7-26.5	4	3.5	00.0-07.8
Some Post-H.S.	156	15,142	23	14.7	08.4-21.0	90	53.8	44.6-63.0	25	21.4	12.8-30.0	15	8.2	03.7-12.7	3	1.9	00.0-04.3
College Graduate	106	6,154	19	18.8	09.4-28.2	64	61.8	50.8-72.8	13	11.5	05.0-18.0	9	7.1	02.4-11.8	1	0.9	00.0-02.7
Less than \$15,000	61	5,687	12	18.6	05.5-31.7	32	57.1	41.8-72.4	7	8.5	02.0-15.0	7	9.8	02.2-17.4	3	6.0	00.0-13.1
\$15,000- 24,999	91	7,098	15	17.4	07.4-27.4	52	49.8	36.7-62.9	14	18.2	06.8-29.6	10	14.6	04.6-24.6			
\$25,000- 34,999	76	5,967	12	14.4	05.6-23.2	41	58.3	45.4-71.2	13	15.0	05.8-24.2	10	12.3	04.1-20.5			
\$35,000- 49,999	65	5,354	11	15.3	05.5-25.1	42	65.4	51.1-79.7	7	10.5	00.3-20.7	2	2.4	00.0-05.7	3	6.5	00.0-15.9
\$50,000- 74,999	57	5,317	6	10.4	01.8-19.0	36	53.3	37.4-69.2	9	27.4	11.3-43.5	5	7.7	00.0-16.3	1	1.3	00.0-03.8
\$75,000+	57	4,546	6	9.9	00.7-19.1	29	44.1	29.2-59.0	11	21.1	08.4-33.8	10	23.7	09.0-38.4	1	1.2	00.0-03.6

^{&#}x27;% = Percentage, C.I.(95%) = Confidence Interval (at 95 percent probability level). Percentages are weighted to population characteristics.

Rows with less than 50 observations have been suppressed.

MODULE 2: DIABETES

TABLE 23: WHEN WAS THE LAST TIME YOU HAD AN EYE EXAM IN WHICH THE PUPILS WERE DILATED? THIS WOULD HAVE MADE YOU TEMPORARILY SENSITIVE TO BRIGHT LIGHT. (EYEEXAM)

Denominator excludes: Respondents who are female or respondents with do not know/refused or with missing responses

Denominator excludes: Respondents that reported not being diagnosed with diabetes

Denominator excludes: Women respondents that reported being diagnosed with diabetes only during pregnancy

Data was weighted using _LANDWT.

DEMOGRAPHIC RESPONDENT NUMBER < 1 Month Ago						1 Month but < 12 Months 1+ Yrs but < 2 Years				2 or more Years				Never			
DEWICGRAPHIC	KESPUNDE	NI NUMBER	<	I WONE	II Ago	i Mont	יו טענ <	12 WOUTENS	1+ 1	5 DUI 4	z redrs	2 (и шоге	i leais		Nev	<i>3</i> 1
GROUPS	TOTAL	WEIGHTED	N	%	C.I. (95%)	N	%	C.I. (95%)	N	%	C.I. (95%)	N	%	C.I. (95%)	N	%	C.I. (95%)
Male	217	20,552	30	12.4	07.5-17.3	120	52.8	45.0-60.6	30	16.0	09.7-22.3	29	14.1	08.4-19.8	8	4.6	01.1-08.1
Non-Native	201	18,964	28	12.7	07.6-17.8	109	51.0	42.8-59.2	28	16.3	09.6-23.0	28	15.0	08.9-21.1	8	5.0	01.1-08.9
55-64	69	5,999	13	19.4	08.4-30.4	35	49.4	35.9-62.9	7	9.4	02.5-16.3	11	18.0	06.6-29.4	3	3.8	00.0-08.3
65+	114	9,101	16	13.2	06.7-19.7	69	62.7	52.9-72.5	16	13.9	06.3-21.5	11	8.0	03.1-12.9	2	2.1	00.0-05.0
H.S. or G.E.D.	66	5,939	9	11.0	03.7-18.3	26	37.4	24.3-50.5	15	19.5	09.3-29.7	14	25.9	12.6-39.2	2	6.2	00.0-15.0
Some Post-H.S.	64	7,097	12	17.5	07.1-27.9	39	56.0	41.7-70.3	6	16.8	03.7-29.9	4	5.6	00.0-11.3	3	4.1	00.0-09.2
College Graduate	56	3,735	9	17.5	05.3-29.7	34	61.0	46.1-75.9	6	10.9	02.1-19.7	6	9.2	01.8-16.6	1	1.4	00.0-04.1

^{&#}x27;% = Percentage, C.I.(95%) = Confidence Interval (at 95 percent probability level). Percentages are weighted to population characteristics.

Rows with less than 50 observations have been suppressed.

MODULE 2: DIABETES

TABLE 24: WHEN WAS THE LAST TIME YOU HAD AN EYE EXAM IN WHICH THE PUPILS WERE DILATED? THIS WOULD HAVE MADE YOU TEMPORARILY SENSITIVE TO BRIGHT LIGHT. (EYEEXAM)

 $\label{lem:percentage} \textbf{Denominator excludes: Respondents who are male or respondents with do not know/refused or with missing responses}$

Denominator excludes: Respondents that reported not being diagnosed with diabetes

Denominator excludes: Women respondents that reported being diagnosed with diabetes only during pregnancy

Data was weighted using _LANDWT.

DEMOGRAPHIC	RESPONDE	NT NUMBER	<	1 Mont	h Ago	1 Mont	h but <	12 Months	1+ Y	rs but •	< 2 Years	2 (or more	Years		Nev	er
GROUPS	TOTAL	WEIGHTED	N	%	C.I. (95%)	N	%	C.I. (95%)	N	%	C.I. (95%)	N	%	C.I. (95%)	N	%	C.I. (95%)
Female	281	19,951	49	16.9	11.2-22.6	157	55.3	48.0-62.6	44	17.4	11.5-23.3	29	10.0	05.9-14.1	2	0.5	00.0-01.1
Non-Native	262	17,368	43	14.0	09.5-18.5	145	54.5	47.2-61.8	44	19.9	13.4-26.4	28	11.0	06.5-15.5	2	0.5	00.0-01.3
55-64	69	5,189	7	12.2	02.0-22.4	41	61.9	48.0-75.8	14	18.9	08.7-29.1	7	7.1	01.4-12.8			
65+	164	9,156	34	18.2	12.1-24.3	93	58.7	50.3-67.1	20	13.2	06.9-19.5	15	8.9	04.2-13.6	2	1.0	00.0-02.4
H.S. or G.E.D.	112	7,178	23	22.3	11.3-33.3	58	50.4	39.0-61.8	15	12.5	05.6-19.4	14	13.5	05.7-21.3	2	1.3	00.0-03.1
Some Post-H.S.	92	8,045	11	12.1	04.5-19.7	51	52.0	39.8-64.2	19	25.4	13.8-37.0	11	10.5	03.8-17.2			
College Graduate	50	2,420	10	20.6	05.7-35.5	30	62.9	46.8-79.0	7	12.6	03.0-22.2	3	3.9	00.0-08.4			
\$15,000- 24,999	62	4.184	10	17.3	05.1-29.5	35	47.8	31.5-64.1	10	21.6	05.9-37.3	7	13.4	03.0-23.8			

^{&#}x27;% = Percentage, C.I.(95%) = Confidence Interval (at 95 percent probability level). Percentages are weighted to population characteristics.

Rows with less than 50 observations have been suppressed.

MODULE 2: DIABETES

TABLE 25: HAS A DOCTOR EVER TOLD YOU THAT DIABETES HAS AFFECTED YOUR EYES OR THAT YOU HAD RETINOPATHY? (DIABEYE)

Denominator excludes: Respondents with do not know/refused/missing responses Denominator excludes: Respondents that reported not being diagnosed with diabetes

Denominator excludes: Women respondents that reported being diagnosed with diabetes only during pregnancy

Data was weighted using _LANDWT.

	Data W	ab weightee	a abii	.9						
DEMOGRAPHIC	RESPONDE	NT NUMBER		Yes	3		No			
GROUPS	TOTAL	WEIGHTED	N	%	C.I. (95%)	N	%	C.I. (95%)		
TOTAL	499	40,506	90	18.9	14.8-23.0	409	81.1	77.0-85.2		
Male	218	20,800	46	22.1	15.6-28.6	172	77.9	71.4-84.4		
Female	281	19,706	44	15.5	10.4-20.6	237	84.5	79.4-89.6		
Non-Native	464	36,685	82	18.5	14.2-22.8	382	81.5	77.2-85.8		
45-54	63	7,678	13	24.1	11.4-36.8	50	75.9	63.2-88.6		
55-64	139	10,957	22	16.7	09.6-23.8	117	83.3	76.2-90.4		
65+	277	18,415	53	19.8	14.5-25.1	224	80.2	74.9-85.		
Less Than H.S.	55	5,721	12	20.5	08.7-32.3	43	79.5	67.7-91.		
H.S. or G.E.D.	178	13,218	34	18.9	12.4-25.4	144	81.1	74.6-87.		
Some Post-H.S.	157	15,264	29	19.8	12.4-27.2	128	80.2	72.8-87.		
College Graduate	107	6,230	14	14.3	04.7-23.9	93	85.7	76.1-95.		
Less than \$15,000	61	5,482	22	34.0	20.5-47.5	39	66.0	52.5-79.		
\$15,000- 24,999	93	7,361	17	20.7	09.9-31.5	76	79.3	68.5-90.		
\$25,000- 34,999	75	5,934	16	24.4	12.2-36.6	59	75.6	63.4-87.		
\$35,000- 49,999	65	5,354	11	16.2	04.8-27.6	54	83.8	72.4-95.		
\$50,000- 74,999	58	5,385	6	12.5	02.3-22.7	52	87.5	77.3-97.		
\$75,000+	57	4,546	6	10.3	01.9-18.7	51	89.7	81.3-98.		

NOTE:

^{&#}x27;% = Percentage, C.I.(95%) = Confidence Interval (at 95 percent probability level). Percentages are weighted to population characteristics. Rows with less than 50 observations have been suppressed.

MODULE 2: DIABETES

TABLE 26: HAS A DOCTOR EVER TOLD YOU THAT DIABETES HAS AFFECTED YOUR EYES OR THAT YOU HAD RETINOPATHY? (DIABEYE)

Denominator excludes: Respondents who are female or respondents with do not know/refused or with missing responses Denominator excludes: Respondents that reported not being diagnosed with diabetes

Denominator excludes: Women respondents that reported being diagnosed with diabetes only during pregnancy

Data was weighted using _LANDWT.

DEMOGRAPHIC	RESPONDE		Yes	3	No			
GROUPS	TOTAL	WEIGHTED	N	%	C.I. (95%)	N	%	C.I. (95%)
Male	218	20,800	46	22.1	15.6-28.6	172	77.9	71.4-84.4
Non-Native	202	19,212	42	21.8	14.9-28.7	160	78.2	71.3-85.1
55-64	69	5,999	11	15.2	06.2-24.2	58	84.8	75.8-93.8
65+	114	9,274	28	24.9	16.1-33.7	86	75.1	66.3-83.9
H.S. or G.E.D.	66	5,953	12	19.9	09.1-30.7	54	80.1	69.3-90.9
Some Post-H.S.	63	7,039	14	22.7	10.9-34.5	49	77.3	65.5-89.1
College Graduate	57	3,810	10	19.3	04.8-33.8	47	80.7	66.2-95.2

^{&#}x27;% = Percentage, C.I.(95%) = Confidence Interval (at 95 percent probability level). Percentages are weighted to population characteristics. Rows with less than 50 observations have been suppressed.

MODULE 2: DIABETES

TABLE 27: HAS A DOCTOR EVER TOLD YOU THAT DIABETES HAS AFFECTED YOUR EYES OR THAT YOU HAD RETINOPATHY? (DIABEYE)

Denominator excludes: Respondents who are male or respondents with do not know/refused or with missing responses Denominator excludes: Respondents that reported not being diagnosed with diabetes

Denominator excludes: Women respondents that reported being diagnosed with diabetes only during pregnancy

Data was weighted using _LANDWT.

DEMOGRAPHIC	RESPONDE	RESPONDENT NUMBER			s	No			
GROUPS	TOTAL	WEIGHTED	N	%	C.I. (95%)	N	%	C.I. (95%)	
Female	281	19,706	44	15.5	10.4-20.6	237	84.5	79.4-89.6	
Non-Native	262	17,473	40	14.9	09.8-20.0	222	85.1	80.0-90.2	
55-64	70	4,958	11	18.4	07.0-29.8	59	81.6	70.2-93.0	
65+	163	9,141	25	14.7	08.8-20.6	138	85.3	79.4-91.2	
H.S. or G.E.D.	112	7,264	22	18.2	10.4-26.0	90	81.8	74.0-89.6	
Some Post-H.S.	94	8,224	15	17.4	07.8-27.0	79	82.6	73.0-92.2	
College Graduate	50	2,420	4	6.4	00.0-12.9	46	93.6	87.1-0100	
\$15,000- 24,999	64	4,412	10	20.1	05.4-34.8	54	79.9	65.2-94.6	

^{&#}x27;% = Percentage, C.I.(95%) = Confidence Interval (at 95 percent probability level). Percentages are weighted to population characteristics. Rows with less than 50 observations have been suppressed.

MODULE 2: DIABETES

TABLE 28: HAVE YOU EVER TAKEN A COURSE OR CLASS IN HOW TO MANAGE YOUR DIABETES YOURSELF? (DIABEDU)

Denominator excludes: Respondents with do not know/refused/missing responses Denominator excludes: Respondents that reported not being diagnosed with diabetes Denominator excludes: Women respondents that reported being diagnosed with diabetes only during pregnancy

Data was weighted using _LANDWT.

	Data W	ab wergiice	a abiii		TIDNI .					
DEMOGRAPHIC	RESPONDE	NT NUMBER		Yes	3		No			
GROUPS	TOTAL	WEIGHTED	N	%	C.I. (95%)	N	%	C.I. (95%)		
TOTAL	502	41,003	314	62.4	57.3-67.5	188	37.6	32.5-42.7		
Male	217	20,656	126	57.1	49.5-64.7	91	42.9	35.3-50.5		
Female	285	20,347	188	67.7	61.2-74.2	97	32.3	25.8-38.8		
Non-Native	467	36,818	293	61.9	56.6-67.2	174	38.1	32.8-43.4		
45-54	63	7,678	48	74.7	62.5-86.9	15	25.3	13.1-37.5		
55-64	139	11,245	91	61.3	51.7-70.9	48	38.7	29.1-48.3		
65+	280	18,624	161	56.1	49.4-62.8	119	43.9	37.2-50.6		
Less Than H.S.	57	6,232	24	46.1	30.8-61.4	33	53.9	38.6-69.2		
H.S. or G.E.D.	180	13,293	104	60.2	51.8-68.6	76	39.8	31.4-48.2		
Some Post-H.S.	156	15,174	106	67.7	59.1-76.3	50	32.3	23.7-40.9		
College Graduate	107	6,230	78	69.8	59.6-80.0	29	30.2	20.0-40.4		
Less than \$15,000	62	5,903	43	70.3	57.0-83.6	19	29.7	16.4-43.0		
\$15,000- 24,999	94	7,419	61	61.0	48.3-73.7	33	39.0	26.3-51.7		
\$25,000- 34,999	76	5,967	52	75.0	64.8-85.2	24	25.0	14.8-35.2		
\$35,000-49,999	64	5,220	30	47.6	33.1-62.1	34	52.4	37.9-66.9		
\$50,000-74,999	58	5,407	39	63.2	48.3-78.1	19	36.8	21.9-51.7		
\$75,000+	56	4,456	36	63.4	48.5-78.3	20	36.6	21.7-51.5		

NOTE:

^{&#}x27;% = Percentage, C.I.(95%) = Confidence Interval (at 95 percent probability level). Percentages are weighted to population characteristics. Rows with less than 50 observations have been suppressed.

MODULE 2: DIABETES

TABLE 29: HAVE YOU EVER TAKEN A COURSE OR CLASS IN HOW TO MANAGE YOUR DIABETES YOURSELF? (DIABEDU)

Denominator excludes: Respondents who are female or respondents with do not know/refused or with missing responses Denominator excludes: Respondents that reported not being diagnosed with diabetes

Denominator excludes: Women respondents that reported being diagnosed with diabetes only during pregnancy

Data was weighted using _LANDWT.

DEMOGRAPHIC	RESPONDE	RESPONDENT NUMBER			3	No			
GROUPS	TOTAL	WEIGHTED	N	%	C.I. (95%)	N	%	C.I. (95%)	
Male	217	20,656	126	57.1	49.5-64.7	91	42.9	35.3-50.5	
Non-Native	202	19,125	117	56.2	48.2-64.2	85	43.8	35.8-51.8	
55-64	68	5,865	42	55.2	41.5-68.9	26	44.8	31.1-58.5	
65+	114	9,263	59	47.8	37.6-58.0	55	52.2	42.0-62.4	
H.S. or G.E.D.	66	5,898	36	58.6	45.3-71.9	30	41.4	28.1-54.7	
Some Post-H.S.	62	6,950	39	61.9	48.2-75.6	23	38.1	24.4-51.8	
College Graduate	57	3,810	38	64.2	49.7-78.7	19	35.8	21.3-50.3	

^{&#}x27;% = Percentage, C.I.(95%) = Confidence Interval (at 95 percent probability level). Percentages are weighted to population characteristics. Rows with less than 50 observations have been suppressed.

MODULE 2: DIABETES

TABLE 30: HAVE YOU EVER TAKEN A COURSE OR CLASS IN HOW TO MANAGE YOUR DIABETES YOURSELF? (DIABEDU)

Denominator excludes: Respondents who are male or respondents with do not know/refused or with missing responses Denominator excludes: Respondents that reported not being diagnosed with diabetes

Denominator excludes: Women respondents that reported being diagnosed with diabetes only during pregnancy

Data was weighted using _LANDWT.

	LIDNII.								
DEMOGRAPHIC	RESPONDE	RESPONDENT NUMBER			3	No			
GROUPS	TOTAL	WEIGHTED	N	%	C.I. (95%)	N	%	C.I. (95%)	
Female	285	20,347	188	67.7	61.0-74.4	97	32.3	25.6-39.0	
Non-Native	265	17,693	176	68.1	61.4-74.8	89	31.9	25.2-38.6	
55-64	71	5,379	49	68.0	55.3-80.7	22	32.0	19.3-44.7	
65+	166	9,361	102	64.3	56.5-72.1	64	35.7	27.9-43.5	
H.S. or G.E.D.	114	7,395	68	61.4	50.6-72.2	46	38.6	27.8-49.4	
Some Post-H.S.	94	8,224	67	72.6	61.6-83.6	27	27.4	16.4-38.4	
College Graduate	50	2,420	40	78.6	65.5-91.7	10	21.4	08.3-34.5	
\$15,000- 24,999	64	4,412	44	69.8	54.7-84.9	20	30.2	15.1-45.3	

^{&#}x27;% = Percentage, C.I.(95%) = Confidence Interval (at 95 percent probability level). Percentages are weighted to population characteristics. Rows with less than 50 observations have been suppressed.

MODULE 10: ACTIONS TO CONTROL HIGH BLOOD PRESSURE CONTENTS

	EN THE DENOMINATOR IS NOT BASED ON THE TOTAL POPULATION SURVEYED, THE TITLE CONTAINS A DESCRIPTION OF WHAT POPULAT EXCLUDED FROM THE DENOMINATOR. IF THE TITLE DOES NOT DESCRIBE THE DENOMINATOR, THE TABLE IS BASED ON ALL RESPONDEN TO THE STATES SURVEY.	TS
* * * * *	* * * * * * * * * * * * * * * * * * *	
TABLE	ARE YOU NOW DOING ANY OF THE FOLLOWING TO HELP LOWER OR CONTROL YOUR HIGH BLOOD PRESSURE? (ARE YOU) CHANGING	PAGE
	YOUR EATING HABITS (TO HELP LOWER OR CONTROL YOUR HIGH BLOOD PRESSURE)? (BPEATHBT)	
1	MALES AND FEMALES	36
2	MALES ONLY	37
3	FEMALES ONLY	38
	[ARE YOU NOW DOING ANY OF THE FOLLOWING TO HELP LOWER OR CONTROL YOUR HIGH BLOOD PRESSURE?] (ARE YOU) CUTTING DOWN ON SALT (TO HELP LOWER OR CONTROL YOUR HIGH BLOOD PRESSURE)? (BPSALT)	
4	MALES AND FEMALES	39
5	MALES ONLY	40
6	FEMALES ONLY	41
	[ARE YOU NOW DOING ANY OF THE FOLLOWING TO HELP LOWER OR CONTROL YOUR HIGH BLOOD PRESSURE?] (ARE YOU) REDUCING ALCOHOL USE (TO HELP LOWER OR CONTROL YOUR HIGH BLOOD PRESSURE)? (BPALCHOL)	
7	MALES AND FEMALES	42
8	MALES ONLY	43
9	FEMALES ONLY	44
	[ARE YOU NOW DOING ANY OF THE FOLLOWING TO HELP LOWER OR CONTROL YOUR HIGH BLOOD PRESSURE?] (ARE YOU) EXERCISING (TO HELP LOWER OR CONTROL YOUR HIGH BLOOD PRESSURE)? (BPEXER)	
10	MALES AND FEMALES	45
11	MALES ONLY	46
12	FEMALES ONLY	47
	HAS A DOCTOR OR OTHER HEALTH PROFESSIONAL EVER ADVISED YOU TO DO ANY OF THE FOLLOWING TO HELP LOWER OR CONTROL YOUR HIGH BLOOD PRESSURE? (EVER ADVISED YOU TO) CHANGE YOUR EATING HABITS (TO HELP LOWER OR CONTROL YOUR HIGH BLOOD PRESSURE)? (BPEATADV)	
13	MALES AND FEMALES	48
14	MALES ONLY	49
15	FEMALES ONLY	50
	[HAS A DOCTOR OR OTHER HEALTH PROFESSIONAL EVER ADVISED YOU TO DO ANY OF THE FOLLOWING TO HELP LOWER OR CONTROL YOUR HIGH BLOOD PRESSURE?] (EVER ADVISED YOU TO) CUT DOWN ON SALT (TO HELP LOWER OR CONTROL YOUR HIGH BLOOD PRESSURE)? (BPSLTADV)	
16	MALES AND FEMALES	51
17	MALES ONLY	52
18	FEMALES ONLY	53

MODULE 10: ACTIONS TO CONTROL HIGH BLOOD PRESSURE CONTENTS

	MODULE 10. MCHOIG TO CONTROL MIGHT BEOOD I RESSURE CONTENTS	
* NOTE	3 * * * * * * * * * * * * * * * * * * *	* * * *
WHE	EN THE DENOMINATOR IS NOT BASED ON THE TOTAL POPULATION SURVEYED, THE TITLE CONTAINS A DESCRIPTION OF WHAT POPULAT	ION
IS E	EXCLUDED FROM THE DENOMINATOR. IF THE TITLE DOES NOT DESCRIBE THE DENOMINATOR, THE TABLE IS BASED ON ALL RESPONDEN	TS
	TO THE STATES SURVEY.	
* * * *	***************************************	* * *
TABLE		PAGE
	[HAS A DOCTOR OR OTHER HEALTH PROFESSIONAL EVER ADVISED YOU TO DO ANY OF THE FOLLOWING TO HELP LOWER OR CONTROL	
	YOUR HIGH BLOOD PRESSURE?] (EVER ADVISED YOU TO) REDUCE ALCOHOL USE (TO HELP LOWER OR CONTROL YOUR HIGH BLOOD	
	PRESSURE)? (BPALCADV)	
19	MALES AND FEMALES	54
20	MALES ONLY	55
21	FEMALES ONLY	56

MODULE 10: ACTIONS TO CONTROL HIGH BLOOD PRESSURE CONTENTS

* NOTE	, *************************************	* * * *
	EN THE DENOMINATOR IS NOT BASED ON THE TOTAL POPULATION SURVEYED, THE TITLE CONTAINS A DESCRIPTION OF WHAT POPULAT EXCLUDED FROM THE DENOMINATOR. IF THE TITLE DOES NOT DESCRIBE THE DENOMINATOR, THE TABLE IS BASED ON ALL RESPONDEN TO THE STATES SURVEY.	
* * * *	*******************************	* * *
TABLE		PAGE
	[HAS A DOCTOR OR OTHER HEALTH PROFESSIONAL EVER ADVISED YOU TO DO ANY OF THE FOLLOWING TO HELP LOWER OR CONTROL YOUR HIGH BLOOD PRESSURE?] (EVER ADVISED YOU TO) EXERCISE (TO HELP LOWER OR CONTROL YOUR HIGH BLOOD PRESSURE)? (BPEXRADV)	
22	MALES AND FEMALES	57
23	MALES ONLY	58
24	FEMALES ONLY	59
	[HAS A DOCTOR OR OTHER HEALTH PROFESSIONAL EVER ADVISED YOU TO DO ANY OF THE FOLLOWING TO HELP LOWER OR CONTROL YOUR HIGH BLOOD PRESSURE?] (EVER ADVISED YOU TO) TAKE MEDICATION (TO HELP LOWER OR CONTROL YOUR HIGH BLOOD PRESSURE)? (BPMEDADV)	
25	MALES AND FEMALES	60
26	MALES ONLY	61
27	FEMALES ONLY	62
	WERE YOU TOLD ON TWO OR MORE DIFFERENT VISITS TO A DOCTOR OR OTHER HEALTH PROFESSIONAL THAT YOU HAD HIGH BLOOD PRESSURE? (BPHI2MR)	
28	MALES AND FEMALES	63
29	MALES ONLY	64
30	FEMALES ONLY	65

TABLE 1: ARE YOU NOW DOING ANY OF THE FOLLOWING TO HELP LOWER OR CONTROL YOUR HIGH BLOOD PRESSURE? (ARE YOU) CHANGING YOUR EATING HABITS (TO HELP LOWER OR CONTROL YOUR HIGH BLOOD PRESSURE)? (BPEATHBT)

Denominator excludes: Respondents with refused/missing responses

	basa was weighted asing _mmswi.									
DEMOGRAPHIC	RESPONDE	NT NUMBER		Yes	3	No				
GROUPS	TOTAL	WEIGHTED	N	%	C.I. (95%)	N	%	C.I. (95%)		
TOTAL	1598	129,617	1178	73.4	70.5-76.3	420	26.6	23.7-29.5		
Male	640	67,394	443	69.4	64.7-74.1	197	30.6	25.9-35.3		
Female	958	62,223	735	77.7	74.6-80.8	223	22.3	19.2-25.4		
Non-Native	1547	124,620	1139	73.0	70.1-75.9	408	27.0	24.1-29.9		
35-44	77	12,867	60	73.3	60.0-86.6	17	26.7	13.4-40.0		
45-54	210	22,695	157	75.0	68.1-81.9	53	25.0	18.1-31.9		
55-64	434	30,759	323	73.3	68.4-78.2	111	26.7	21.8-31.6		
65+	819	50,690	598	72.2	68.7-75.7	221	27.8	24.3-31.3		
Less Than H.S.	139	14,867	94	64.3	54.5-74.1	45	35.7	25.9-45.5		
H.S. or G.E.D.	532	39,649	386	73.7	69.0-78.4	146	26.3	21.6-31.0		
Some Post-H.S.	482	48,417	354	72.4	66.9-77.9	128	27.6	22.1-33.1		
College Graduate	441	26,259	341	79.4	75.1-83.7	100	20.6	16.3-24.9		
Less than \$15,000	143	12,267	103	77.3	70.0-84.6	40	22.7	15.4-30.0		
\$15,000- 24,999	252	19,720	185	70.2	62.0-78.4	67	29.8	21.6-38.0		
\$25,000- 34,999	197	15,533	143	73.2	64.6-81.8	54	26.8	18.2-35.4		
\$35,000- 49,999	234	18,091	172	71.7	64.4-79.0	62	28.3	21.0-35.6		
\$50,000- 74,999	217	19,959	155	68.6	59.8-77.4	62	31.4	22.6-40.2		
\$75,000+	280	23,769	213	76.8	71.1-82.5	67	23.2	17.5-28.9		

NOTE:

^{&#}x27;% = Percentage, C.I.(95%) = Confidence Interval (at 95 percent probability level). Percentages are weighted to population characteristics.

Rows with less than 50 observations have been suppressed.

TABLE 2: ARE YOU NOW DOING ANY OF THE FOLLOWING TO HELP LOWER OR CONTROL YOUR HIGH BLOOD PRESSURE? (ARE YOU) CHANGING YOUR EATING HABITS (TO HELP LOWER OR CONTROL YOUR HIGH BLOOD PRESSURE)? (BPEATHBT)

Denominator excludes: Respondents who are female or respondents with refused or with missing responses

	9								
DEMOGRAPHIC	RESPONDE	ESPONDENT NUMBER		Yes	3	No			
GROUPS	TOTAL	WEIGHTED	N	%	C.I. (95%)	N	%	C.I. (95%)	
Male	640	67,394	443	69.4	64.7-74.1	197	30.6	25.9-35.3	
Non-Native	618	64,950	425	68.8	63.9-73.7	193	31.2	26.3-36.1	
45-54	98	12,806	66	68.7	58.1-79.3	32	31.3	20.7-41.9	
55-64	201	16,796	139	67.9	60.5-75.3	62	32.1	24.7-39.5	
65+	271	21,306	183	66.3	60.0-72.6	88	33.7	27.4-40.0	
Less Than H.S.	62	8,203	37	56.0	41.5-70.5	25	44.0	29.5-58.5	
H.S. or G.E.D.	203	19,975	134	70.9	63.5-78.3	69	29.1	21.7-36.5	
Some Post-H.S.	181	25,083	123	66.3	57.1-75.5	58	33.7	24.5-42.9	
College Graduate	192	13,789	147	80.0	73.9-86.1	45	20.0	13.9-26.1	
\$15,000- 24,999	70	7,564	48	59.8	43.3-76.3	22	40.2	23.7-56.7	
\$25,000- 34,999	82	9,474	55	71.7	58.4-85.0	27	28.3	15.0-41.6	
\$35,000-49,999	114	10,619	76	66.1	56.1-76.1	38	33.9	23.9-43.9	
\$50,000-74,999	113	13,063	75	63.1	50.8-75.4	38	36.9	24.6-49.2	
\$75,000+	159	15,444	118	76.4	69.1-83.7	41	23.6	16.3-30.9	

NOTE:

^{&#}x27;% = Percentage, C.I.(95%) = Confidence Interval (at 95 percent probability level). Percentages are weighted to population characteristics.

Rows with less than 50 observations have been suppressed.

TABLE 3: ARE YOU NOW DOING ANY OF THE FOLLOWING TO HELP LOWER OR CONTROL YOUR HIGH BLOOD PRESSURE? (ARE YOU) CHANGING YOUR EATING HABITS (TO HELP LOWER OR CONTROL YOUR HIGH BLOOD PRESSURE)? (BPEATHBT)

Denominator excludes: Respondents who are male or respondents with refused or with missing responses

Data was weighted using _LANDWT.

DEMOGRAPHIC	RESPONDE		Yes	3	No					
GROUPS	TOTAL	WEIGHTED	N	%	C.I. (95%)	N	%	C.I. (95%)		
Female	958	62,223	735	77.7	74.6-80.8	223	22.3	19.2-25.4		
Non-Native	929	59,670	714	77.5	74.4-80.6	215	22.5	19.4-25.6		
45-54	112	9,890	91	83.1	75.8-90.4	21	16.9	09.6-24.2		
55-64	233	13,962	184	79.8	74.3-85.3	49	20.2	14.7-25.7		
65+	548	29,384	415	76.4	72.7-80.1	133	23.6	19.9-27.3		
Less Than H.S.	77	6,663	57	74.4	62.8-86.0	20	25.6	14.0-37.2		
H.S. or G.E.D.	329	19,673	252	76.6	70.9-82.3	77	23.4	17.7-29.1		
Some Post-H.S.	301	23,334	231	79.0	74.1-83.9	70	21.0	16.1-25.9		
College Graduate	249	12,470	194	78.8	72.9-84.7	55	21.2	15.3-27.1		
Less than \$15,000	103	7,663	74	76.3	67.5-85.1	29	23.7	14.9-32.5		
\$15,000- 24,999	182	12,156	137	76.7	69.8-83.6	45	23.3	16.4-30.2		
\$25,000- 34,999	115	6,059	88	75.7	67.1-84.3	27	24.3	15.7-32.9		
\$35,000-49,999	120	7,472	96	79.7	69.3-90.1	24	20.3	09.9-30.7		
\$50,000-74,999	104	6,895	80	79.2	70.8-87.6	24	20.8	12.4-29.2		
\$75,000+	121	8,324	95	77.7	68.9-86.5	26	22.3	13.5-31.1		

NOTE:

July 26, 2012

^{&#}x27;% = Percentage, C.I.(95%) = Confidence Interval (at 95 percent probability level). Percentages are weighted to population characteristics.

Rows with less than 50 observations have been suppressed.

TABLE 4: [ARE YOU NOW DOING ANY OF THE FOLLOWING TO HELP LOWER OR CONTROL YOUR HIGH BLOOD PRESSURE?] (ARE YOU) CUTTING DOWN ON SALT (TO HELP LOWER OR CONTROL YOUR HIGH BLOOD PRESSURE)? (BPSALT)

Denominator excludes: Respondents with refused/missing responses

Data was weighted using _LANDWT.

	ab wei	wergheed abing _mmbmr.									
DEMOGRAPHIC	RESPONDE	NT NUMBER	Yes			No			Do	o not u	se salt
GROUPS	TOTAL	WEIGHTED	N	%	C.I. (95%)	N	%	C.I. (95%)	N	%	C.I. (95%)
TOTAL	1604	130,017	1261	78.1	75.6-80.6	242	15.8	13.4-18.2	101	6.0	04.6-07.4
Male	643	67,727	480	75.4	71.3-79.5	120	18.0	14.3-21.7	43	6.6	04.2-09.0
Female	961	62,290	781	81.0	77.9-84.1	122	13.5	10.8-16.2	58	5.5	03.9-07.1
Non-Native	1553	125,020	1225	78.1	75.4-80.8	233	16.0	13.6-18.4	95	5.9	04.3-07.5
35-44	77	12,867	56	70.5	58.2-82.8	16	21.3	10.3-32.3	5	8.2	00.6-15.8
45-54	211	23,136	170	82.1	76.0-88.2	34	13.9	08.6-19.2	7	4.0	00.7-07.3
55-64	437	30,896	344	78.1	73.6-82.6	72	17.1	13.0-21.2	21	4.8	02.4-07.2
65+	822	50,792	653	79.1	76.0-82.2	105	13.2	10.7-15.7	64	7.6	05.6-09.6
Less Than H.S.	138	14,781	106	71.2	61.4-81.0	24	21.9	12.5-31.3	8	6.8	01.9-11.7
H.S. or G.E.D.	536	40,219	423	78.2	73.5-82.9	70	14.2	10.3-18.1	43	7.6	04.7-10.5
Some Post-H.S.	484	48,284	388	80.5	76.2-84.8	71	14.0	10.3-17.7	25	5.5	03.0-08.0
College Graduate	442	26,308	340	77.2	72.5-81.9	77	18.6	14.3-22.9	25	4.2	02.4-06.0
Less than \$15,000	144	12,290	115	81.7	74.6-88.8	19	10.4	05.5-15.3	10	7.9	02.4-13.4
\$15,000- 24,999	254	19,795	207	80.6	74.5-86.7	32	13.7	08.4-19.0	15	5.7	02.2-09.2
\$25,000- 34,999	196	15,443	155	81.4	74.7-88.1	27	13.7	07.6-19.8	14	4.9	02.2-07.6
\$35,000- 49,999	236	18,199	188	77.2	70.3-84.1	30	15.6	09.5-21.7	18	7.2	03.3-11.1
\$50,000- 74,999	217	19,735	160	71.3	62.9-79.7	44	22.3	14.5-30.1	13	6.4	01.7-11.1
\$75,000+	280	23,811	217	78.7	73.2-84.2	51	16.4	11.7-21.1	12	4.9	01.8-08.0

^{&#}x27;% = Percentage, C.I.(95%) = Confidence Interval (at 95 percent probability level). Percentages are weighted to population characteristics.

Rows with less than 50 observations have been suppressed.

TABLE 5: [ARE YOU NOW DOING ANY OF THE FOLLOWING TO HELP LOWER OR CONTROL YOUR HIGH BLOOD PRESSURE?] (ARE YOU) CUTTING DOWN ON SALT (TO HELP LOWER OR CONTROL YOUR HIGH BLOOD PRESSURE)? (BPSALT)

Denominator excludes: Respondents who are female or respondents with refused or with missing responses

Data was weighted using _LANDWT.

DEMOGRAPHIC	RESPONDE	RESPONDENT NUMBER		Yes		No			Do not use salt		
GROUPS	TOTAL	WEIGHTED	N	%	C.I. (95%)	N	%	C.I. (95%)	N	%	C.I. (95%)
Male	643	67,727	480	75.4	71.3-79.5	120	18.0	14.3-21.7	43	6.6	04.2-09.0
Non-Native	621	65,283	464	75.4	71.1-79.7	115	18.0	14.3-21.7	42	6.6	04.2-09.0
45-54	100	13,335	76	79.1	70.1-88.1	20	15.7	07.9-23.5	4	5.2	00.0-10.7
55-64	202	16,834	149	73.2	66.3-80.1	43	21.9	15.4-28.4	10	4.9	01.4-08.4
65+	272	21,350	203	74.4	68.7-80.1	44	16.4	11.5-21.3	25	9.2	05.5-12.9
Less Than H.S.	62	8,203	40	61.2	46.7-75.7	17	29.8	15.7-43.9	5	9.1	01.1-17.1
H.S. or G.E.D.	205	20,456	152	75.1	67.5-82.7	34	15.8	09.7-21.9	19	9.1	04.0-14.2
Some Post-H.S.	180	24,846	140	79.4	72.5-86.3	30	14.9	09.0-20.8	10	5.7	01.8-09.6
College Graduate	194	13,879	146	76.8	70.1-83.5	39	20.2	13.7-26.7	9	3.0	01.0-05.0
\$15,000- 24,999	70	7,564	53	74.1	61.6-86.6	12	20.4	08.6-32.2	5	5.5	00.2-10.8
\$25,000- 34,999	81	9,384	65	85.6	77.2-94.0	10	10.1	02.7-17.5	6	4.3	00.8-07.8
\$35,000- 49,999	115	10,671	80	70.6	61.2-80.0	24	20.3	12.3-28.3	11	9.1	03.0-15.2
\$50,000- 74,999	113	12,823	81	68.0	56.0-80.0	24	24.1	12.9-35.3	8	7.9	00.8-15.0
\$75.000+	160	15.575	124	80.4	73.5-87.3	29	14.6	09.1-20.1	7	5.1	00.8-09.4

^{&#}x27;% = Percentage, C.I.(95%) = Confidence Interval (at 95 percent probability level). Percentages are weighted to population characteristics.

Rows with less than 50 observations have been suppressed.

TABLE 6: [ARE YOU NOW DOING ANY OF THE FOLLOWING TO HELP LOWER OR CONTROL YOUR HIGH BLOOD PRESSURE?] (ARE YOU) CUTTING DOWN ON SALT (TO HELP LOWER OR CONTROL YOUR HIGH BLOOD PRESSURE)? (BPSALT)

Denominator excludes: Respondents who are male or respondents with refused or with missing responses

Data was weighted using _LANDWT.

DEMOCRABUIC	DECDONDE	NT NUMBER	ib wei	_		INDNI			ъ.		
DEMOGRAPHIC		NT NUMBER		Yes			No				se salt
GROUPS	TOTAL	WEIGHTED	N	%	C.I. (95%)	N	%	C.I. (95%)	N	%	C.I. (95%)
Female	961	62,290	781	81.0	77.9-84.1	122	13.5	10.8-16.2	58	5.5	03.9-07.1
Non-Native	932	59,737	761	81.1	78.0-84.2	118	13.7	10.8-16.6	53	5.2	03.6-06.8
45-54	111	9,801	94	86.3	79.2-93.4	14	11.4	04.7-18.1	3	2.3	00.0-05.0
55-64	235	14,062	195	83.9	79.0-88.8	29	11.3	07.2-15.4	11	4.8	01.9-07.7
65+	550	29,442	450	82.5	79.2-85.8	61	11.0	08.3-13.7	39	6.5	04.3-08.7
Less Than H.S.	76	6,578	66	83.8	73.0-94.6	7	12.2	01.8-22.6	3	4.0	00.0-08.5
H.S. or G.E.D.	331	19,763	271	81.4	75.9-86.9	36	12.5	07.4-17.6	24	6.1	03.6-08.6
Some Post-H.S.	304	23,439	248	81.6	76.5-86.7	41	13.1	08.8-17.4	15	5.3	02.2-08.4
College Graduate	248	12,429	194	77.7	71.4-84.0	38	16.7	11.0-22.4	16	5.6	02.5-08.7
Less than \$15,000	104	7,686	84	84.3	77.0-91.6	13	9.7	04.2-15.2	7	6.0	01.1-10.9
\$15,000- 24,999	184	12,230	154	84.6	78.3-90.9	20	9.6	05.1-14.1	10	5.8	01.1-10.5
\$25,000- 34,999	115	6,059	90	75.0	65.6-84.4	17	19.2	10.2-28.2	8	5.8	01.7-09.9
\$35,000- 49,999	121	7,529	108	86.5	76.3-96.7	6	9.0	00.0-18.8	7	4.5	00.8-08.2
\$50,000-74,999	104	6,912	79	77.5	68.7-86.3	20	18.9	10.5-27.3	5	3.6	00.3-06.9
\$75,000+	120	8,235	93	75.6	66.2-85.0	22	19.8	11.0-28.6	5	4.5	00.4-08.6

^{&#}x27;% = Percentage, C.I.(95%) = Confidence Interval (at 95 percent probability level). Percentages are weighted to population characteristics.

Rows with less than 50 observations have been suppressed.

TABLE 7: [ARE YOU NOW DOING ANY OF THE FOLLOWING TO HELP LOWER OR CONTROL YOUR HIGH BLOOD PRESSURE?] (ARE YOU) REDUCING ALCOHOL USE (TO HELP LOWER OR CONTROL YOUR HIGH BLOOD PRESSURE)? (BPALCHOL)

Denominator excludes: Respondents with refused/missing responses

Data was weighted using _LANDWT.

		Data w	ab wei	_	abing _i	anibiri.					
DEMOGRAPHIC	RESPONDE	NT NUMBER		Yes	i		No			Do not o	drink
GROUPS	TOTAL	WEIGHTED	N	%	C.I. (95%)	N	%	C.I. (95%)	N	%	C.I. (95%)
TOTAL	1598	129,453	612	42.4	39.1-45.7	400	24.8	22.1-27.5	586	32.9	30.0-35.8
Male	638	67,179	269	47.3	42.0-52.6	195	28.4	24.1-32.7	174	24.3	20.0-28.6
Female	960	62,273	343	37.0	33.3-40.7	205	20.9	17.8-24.0	412	42.1	38.4-45.8
Non-Native	1547	124,455	592	42.0	38.7-45.3	391	25.4	22.7-28.1	564	32.6	29.7-35.5
35-44	77	12,867	43	55.7	42.2-69.2	21	23.6	13.0-34.2	13	20.8	08.5-33.1
45-54	209	22,509	96	47.4	39.6-55.2	58	26.2	19.3-33.1	55	26.5	19.4-33.6
55-64	434	30,781	190	45.0	39.7-50.3	124	28.0	23.3-32.7	120	27.0	22.3-31.7
65+	821	50,740	256	31.2	27.7-34.7	181	22.8	19.5-26.1	384	46.0	42.1-49.9
Less Than H.S.	139	14,867	45	37.4	27.6-47.2	28	20.5	12.5-28.5	66	42.2	33.0-51.4
H.S. or G.E.D.	530	39,529	213	44.8	38.7-50.9	105	21.5	16.8-26.2	212	33.7	28.8-38.6
Some Post-H.S.	484	48,374	191	42.0	36.1-47.9	120	24.4	19.7-29.1	173	33.6	28.1-39.1
College Graduate	441	26,257	163	43.0	36.9-49.1	146	32.2	26.9-37.5	132	24.8	20.1-29.5
Less than \$15,000	144	12,290	54	39.0	29.2-48.8	29	22.9	14.3-31.5	61	38.1	28.9-47.3
\$15,000- 24,999	251	19,501	104	40.9	33.1-48.7	41	14.7	09.8-19.6	106	44.5	36.3-52.7
\$25,000- 34,999	197	15,533	71	44.3	33.3-55.3	46	24.0	15.4-32.6	80	31.7	23.5-39.9
\$35,000- 49,999	234	18,095	87	37.6	30.2-45.0	59	28.4	21.0-35.8	88	34.0	26.7-41.3
\$50,000- 74,999	219	20,037	89	43.5	34.7-52.3	70	31.2	23.2-39.2	60	25.3	17.1-33.5
\$75,000+	277	23,643	112	49.0	41.7-56.3	101	32.5	26.2-38.8	64	18.6	13.7-23.5

^{&#}x27;% = Percentage, C.I.(95%) = Confidence Interval (at 95 percent probability level). Percentages are weighted to population characteristics.

Rows with less than 50 observations have been suppressed.

TABLE 8: [ARE YOU NOW DOING ANY OF THE FOLLOWING TO HELP LOWER OR CONTROL YOUR HIGH BLOOD PRESSURE?] (ARE YOU) REDUCING ALCOHOL USE (TO HELP LOWER OR CONTROL YOUR HIGH BLOOD PRESSURE)? (BPALCHOL)

Denominator excludes: Respondents who are female or respondents with refused or with missing responses

Data was weighted using _LANDWT.

				J			-				
DEMOGRAPHIC	RESPONDE	NT NUMBER	Yes				No		Do not drink		
GROUPS	TOTAL	WEIGHTED	N	%	C.I. (95%)	N	%	C.I. (95%)	N	%	C.I. (95%)
Male	638	67,179	269	47.3	42.0-52.6	195	28.4	24.1-32.7	174	24.3	20.0-28.6
Non-Native	616	64,735	259	47.0	41.5-52.5	189	28.9	24.4-33.4	168	24.1	19.8-28.4
45-54	97	12,620	48	50.0	38.6-61.4	29	28.5	18.1-38.9	20	21.5	11.9-31.1
55-64	200	16,748	89	45.4	37.6-53.2	62	30.8	23.7-37.9	49	23.8	17.3-30.3
65+	271	21,325	89	33.3	27.2-39.4	84	31.5	25.2-37.8	98	35.3	29.0-41.6
Less Than H.S.	62	8,203	21	42.8	28.1-57.5	16	23.6	11.8-35.4	25	33.6	21.1-46.1
H.S. or G.E.D.	202	19,943	79	45.2	35.0-55.4	56	27.2	19.4-35.0	67	27.5	20.2-34.8
Some Post-H.S.	180	24,897	88	50.3	40.7-59.9	54	27.4	19.6-35.2	38	22.3	13.7-30.9
College Graduate	192	13,793	81	48.8	40.0-57.6	68	33.5	25.7-41.3	43	17.7	12.2-23.2
\$15,000- 24,999	69	7,378	31	41.5	26.6-56.4	14	16.8	07.4-26.2	24	41.8	25.3-58.3
\$25,000- 34,999	82	9,474	32	48.3	32.2-64.4	23	27.6	14.1-41.1	27	24.0	13.4-34.6
\$35,000- 49,999	113	10,566	40	34.7	24.9-44.5	31	30.7	20.5-40.9	42	34.6	24.4-44.8
\$50,000- 74,999	114	13,102	48	44.3	32.1-56.5	42	34.0	22.8-45.2	24	21.7	10.5-32.9
\$75,000+	157	15,348	74	57.5	48.5-66.5	52	28.3	20.7-35.9	31	14.2	08.9-19.5

^{&#}x27;% = Percentage, C.I.(95%) = Confidence Interval (at 95 percent probability level). Percentages are weighted to population characteristics.

Rows with less than 50 observations have been suppressed.

TABLE 9: [ARE YOU NOW DOING ANY OF THE FOLLOWING TO HELP LOWER OR CONTROL YOUR HIGH BLOOD PRESSURE?] (ARE YOU) REDUCING ALCOHOL USE (TO HELP LOWER OR CONTROL YOUR HIGH BLOOD PRESSURE)? (BPALCHOL)

Denominator excludes: Respondents who are male or respondents with refused or with missing responses

Data was weighted using _LANDWT.

				J							
DEMOGRAPHIC	RESPONDE	NT NUMBER		Yes	5		No		1	Do not	drink
GROUPS	TOTAL	WEIGHTED	N	%	C.I. (95%)	N	%	C.I. (95%)	N	%	C.I. (95%)
Female	960	62,273	343	37.0	33.3-40.7	205	20.9	17.8-24.0	412	42.1	38.4-45.8
Non-Native	931	59,720	333	36.6	32.9-40.3	202	21.6	18.5-24.7	396	41.8	38.1-45.5
45-54	112	9,890	48	43.9	33.5-54.3	29	23.2	14.8-31.6	35	32.8	23.0-42.6
55-64	234	14,033	101	44.5	37.4-51.6	62	24.6	18.7-30.5	71	30.9	24.2-37.6
65+	550	29,415	167	29.6	25.5-33.7	97	16.5	13.2-19.8	286	53.8	49.3-58.3
Less Than H.S.	77	6,663	24	30.7	19.1-42.3	12	16.6	06.0-27.2	41	52.8	40.3-65.3
H.S. or G.E.D.	328	19,586	134	44.3	37.6-51.0	49	15.7	10.4-21.0	145	40.0	33.9-46.1
Some Post-H.S.	304	23,477	103	33.2	27.3-39.1	66	21.3	16.2-26.4	135	45.5	39.0-52.0
College Graduate	249	12,465	82	36.5	28.7-44.3	78	30.7	23.8-37.6	89	32.8	25.4-40.2
Less than \$15,000	104	7,686	39	37.7	26.3-49.1	14	13.6	06.2-21.0	51	48.7	37.1-60.3
\$15,000- 24,999	182	12,122	73	40.5	31.7-49.3	27	13.4	08.1-18.7	82	46.1	37.3-54.9
\$25,000- 34,999	115	6,059	39	38.0	27.6-48.4	23	18.3	10.9-25.7	53	43.7	33.5-53.9
\$35,000- 49,999	121	7,529	47	41.6	30.4-52.8	28	25.3	14.5-36.1	46	33.1	23.3-42.9
\$50,000- 74,999	105	6,936	41	41.9	30.7-53.1	28	26.0	16.8-35.2	36	32.2	21.6-42.8
\$75,000+	120	8,295	38	33.3	22.9-43.7	49	40.1	29.7-50.5	33	26.6	17.4-35.8

^{&#}x27;% = Percentage, C.I.(95%) = Confidence Interval (at 95 percent probability level). Percentages are weighted to population characteristics.

Rows with less than 50 observations have been suppressed.

TABLE 10: [ARE YOU NOW DOING ANY OF THE FOLLOWING TO HELP LOWER OR CONTROL YOUR HIGH BLOOD PRESSURE?] (ARE YOU) EXERCISING (TO HELP LOWER OR CONTROL YOUR HIGH BLOOD PRESSURE)? (BPEXER)

Denominator excludes: Respondents with refused/missing responses

	zada naz nergireda azing _zinzni.									
DEMOGRAPHIC	RESPONDE	NT NUMBER		Yes	3	No				
GROUPS	TOTAL	WEIGHTED	N	%	C.I. (95%)	N	%	C.I. (95%)		
TOTAL	1590	128,834	1059	66.1	63.0-69.2	531	33.9	30.8-37.0		
Male	637	67,072	428	65.5	60.2-70.8	209	34.5	29.2-39.8		
Female	953	61,761	631	66.8	63.3-70.3	322	33.2	29.7-36.7		
Non-Native	1540	123,998	1027	66.0	62.7-69.3	513	34.0	30.7-37.3		
35-44	77	12,867	54	66.3	52.8-79.8	23	33.7	20.2-47.2		
45-54	209	22,644	142	67.8	60.4-75.2	67	32.2	24.8-39.6		
55-64	432	30,516	295	67.7	62.8-72.6	137	32.3	27.4-37.2		
65+	815	50,362	526	63.6	59.9-67.3	289	36.4	32.7-40.1		
Less Than H.S.	136	14,604	81	61.2	51.4-71.0	55	38.8	29.0-48.6		
H.S. or G.E.D.	525	39,093	343	65.1	58.8-71.4	182	34.9	28.6-41.2		
Some Post-H.S.	484	48,461	291	61.8	56.3-67.3	193	38.2	32.7-43.7		
College Graduate	442	26,290	341	77.8	72.9-82.7	101	22.2	17.3-27.1		
Less than \$15,000	142	12,127	82	64.6	55.6-73.6	60	35.4	26.4-44.4		
\$15,000- 24,999	251	19,604	162	68.0	60.9-75.1	89	32.0	24.9-39.1		
\$25,000- 34,999	196	15,499	116	58.1	47.5-68.7	80	41.9	31.3-52.5		
\$35,000-49,999	235	18,095	143	60.8	53.2-68.4	92	39.2	31.6-46.8		
\$50,000-74,999	216	19,786	159	70.6	62.2-79.0	57	29.4	21.0-37.8		
\$75,000+	279	23,773	207	72.0	65.5-78.5	72	28.0	21.5-34.5		

NOTE:

^{&#}x27;% = Percentage, C.I.(95%) = Confidence Interval (at 95 percent probability level). Percentages are weighted to population characteristics.

Rows with less than 50 observations have been suppressed.

TABLE 11: [ARE YOU NOW DOING ANY OF THE FOLLOWING TO HELP LOWER OR CONTROL YOUR HIGH BLOOD PRESSURE?] (ARE YOU) EXERCISING (TO HELP LOWER OR CONTROL YOUR HIGH BLOOD PRESSURE)? (BPEXER)

Denominator excludes: Respondents who are female or respondents with refused or with missing responses

DEMOGRAPHIC	RESPONDENT NUMBER			Yes	6	No			
GROUPS	TOTAL	WEIGHTED	N	%	C.I. (95%)	N	%	C.I. (95%)	
Male	637	67,072	428	65.5	60.2-70.8	209	34.5	29.2-39.8	
Non-Native	616	64,790	413	65.0	59.7-70.3	203	35.0	29.7-40.3	
45-54	98	12,826	67	68.7	58.1-79.3	31	31.3	20.7-41.9	
55-64	200	16,636	132	66.7	59.4-74.0	68	33.3	26.0-40.6	
65+	270	21,285	182	63.5	56.8-70.2	88	36.5	29.8-43.2	
Less Than H.S.	61	8,081	35	57.6	42.9-72.3	26	42.4	27.7-57.1	
H.S. or G.E.D.	200	19,637	133	62.6	52.0-73.2	67	37.4	26.8-48.0	
Some Post-H.S.	182	25,214	107	60.6	51.4-69.8	75	39.4	30.2-48.6	
College Graduate	192	13,797	151	82.2	76.5-87.9	41	17.8	12.1-23.5	
\$15,000- 24,999	68	7,423	49	76.7	65.3-88.1	19	23.3	11.9-34.7	
\$25,000- 34,999	82	9,474	49	57.7	41.6-73.8	33	42.3	26.2-58.4	
\$35,000- 49,999	115	10,671	69	57.3	46.7-67.9	46	42.7	32.1-53.3	
\$50,000-74,999	113	13,029	77	68.4	57.0-79.8	36	31.6	20.2-43.0	
\$75,000+	158	15,448	114	70.3	61.9-78.7	44	29.7	21.3-38.1	

NOTE:

^{&#}x27;% = Percentage, C.I.(95%) = Confidence Interval (at 95 percent probability level). Percentages are weighted to population characteristics.

Rows with less than 50 observations have been suppressed.

TABLE 12: [ARE YOU NOW DOING ANY OF THE FOLLOWING TO HELP LOWER OR CONTROL YOUR HIGH BLOOD PRESSURE?] (ARE YOU) EXERCISING (TO HELP LOWER OR CONTROL YOUR HIGH BLOOD PRESSURE)? (BPEXER)

Denominator excludes: Respondents who are male or respondents with refused or with missing responses

				·					
DEMOGRAPHIC	RESPONDE		Yes	5	No				
GROUPS	TOTAL	WEIGHTED	N	%	C.I. (95%)	N	%	C.I. (95%)	
Female	953	61,761	631	66.8	63.3-70.3	322	33.2	29.7-36.7	
Non-Native	924	59,208	614	67.0	63.5-70.5	310	33.0	29.5-36.5	
45-54	111	9,818	75	66.6	56.6-76.6	36	33.4	23.4-43.4	
55-64	232	13,880	163	68.9	62.2-75.6	69	31.1	24.4-37.8	
65+	545	29,077	344	63.6	59.1-68.1	201	36.4	31.9-40.9	
Less Than H.S.	75	6,523	46	65.7	53.9-77.5	29	34.3	22.5-46.1	
H.S. or G.E.D.	325	19,456	210	67.6	61.7-73.5	115	32.4	26.5-38.3	
Some Post-H.S.	302	23,247	184	63.1	57.0-69.2	118	36.9	30.8-43.0	
College Graduate	250	12,494	190	73.0	65.4-80.6	60	27.0	19.4-34.6	
Less than \$15,000	103	7,646	57	60.2	49.0-71.4	46	39.8	28.6-51.0	
\$15,000- 24,999	183	12,181	113	62.7	54.3-71.1	70	37.3	28.9-45.7	
\$25,000- 34,999	114	6,025	67	58.7	48.7-68.7	47	41.3	31.3-51.3	
\$35,000- 49,999	120	7,424	74	66.0	55.8-76.2	46	34.0	23.8-44.2	
\$50,000-74,999	103	6,756	82	74.9	63.7-86.1	21	25.1	13.9-36.3	
\$75,000+	121	8,324	93	75.0	65.6-84.4	28	25.0	15.6-34.4	

NOTE:

^{&#}x27;% = Percentage, C.I.(95%) = Confidence Interval (at 95 percent probability level). Percentages are weighted to population characteristics.

Rows with less than 50 observations have been suppressed.

TABLE 13: HAS A DOCTOR OR OTHER HEALTH PROFESSIONAL EVER ADVISED YOU TO DO ANY OF THE FOLLOWING TO HELP LOWER OR CONTROL YOUR HIGH BLOOD PRESSURE? (EVER ADVISED YOU TO) CHANGE YOUR EATING HABITS (TO HELP LOWER OR CONTROL YOUR HIGH BLOOD PRESSURE)? (BPEATADV)

Denominator excludes: Respondents with refused/missing responses

Data was weighted using LANDWT.

	MDWI.							
DEMOGRAPHIC	RESPONDE	RESPONDENT NUMBER			3		No	
GROUPS	TOTAL	WEIGHTED	N	%	C.I. (95%)	N	%	C.I. (95%)
TOTAL	1590	128,983	914	59.5	56.4-62.6	676	40.5	37.4-43.6
Male	635	67,190	390	61.4	56.3-66.5	245	38.6	33.5-43.7
Female	955	61,793	524	57.5	53.8-61.2	431	42.5	38.8-46.2
Non-Native	1540	124,079	878	58.7	55.4-62.0	662	41.3	38.0-44.6
35-44	77	12,867	49	60.3	46.8-73.8	28	39.7	26.2-53.2
45-54	212	23,225	143	65.8	58.2-73.4	69	34.2	26.6-41.8
55-64	434	30,704	288	67.5	62.6-72.4	146	32.5	27.6-37.4
65+	811	49,742	403	51.6	47.7-55.5	408	48.4	44.5-52.3
Less Than H.S.	137	14,625	68	53.7	43.9-63.5	69	46.3	36.5-56.1
H.S. or G.E.D.	530	39,768	302	60.2	54.5-65.9	228	39.8	34.1-45.5
Some Post-H.S.	478	47,966	277	59.1	53.2-65.0	201	40.9	35.0-46.8
College Graduate	442	26,292	267	63.3	57.8-68.8	175	36.7	31.2-42.2
Less than \$15,000	142	12,192	79	60.2	50.6-69.8	63	39.8	30.2-49.4
\$15,000- 24,999	253	19,709	142	57.4	49.2-65.6	111	42.6	34.4-50.8
\$25,000- 34,999	195	15,404	105	54.9	44.1-65.7	90	45.1	34.3-55.9
\$35,000-49,999	232	17,824	134	58.8	51.0-66.6	98	41.2	33.4-49.0
\$50,000-74,999	218	19,965	136	63.5	54.7-72.3	82	36.5	27.7-45.3
\$75,000+	278	23,708	170	64.1	57.4-70.8	108	35.9	29.2-42.6

NOTE:

July 26, 2012

^{&#}x27;% = Percentage, C.I.(95%) = Confidence Interval (at 95 percent probability level). Percentages are weighted to population characteristics.

Rows with less than 50 observations have been suppressed.

TABLE 14: HAS A DOCTOR OR OTHER HEALTH PROFESSIONAL EVER ADVISED YOU TO DO ANY OF THE FOLLOWING TO HELP LOWER OR CONTROL YOUR HIGH BLOOD PRESSURE? (EVER ADVISED YOU TO) CHANGE YOUR EATING HABITS (TO HELP LOWER OR CONTROL YOUR HIGH BLOOD PRESSURE)? (BPEATADV)

Denominator excludes: Respondents who are female or respondents with refused or with missing responses

Data was weighted using _LANDWT.

DEMOGRAPHIC	RESPONDE	NT NUMBER		Yes	3	No			
GROUPS	TOTAL	WEIGHTED	N	%	C.I. (95%)	N	%	C.I. (95%)	
Male	635	67,190	390	61.4	56.3-66.5	245	38.6	33.5-43.7	
Non-Native	614	64,839	373	60.6	55.3-65.9	241	39.4	34.1-44.7	
45-54	100	13,335	68	63.5	52.3-74.7	32	36.5	25.3-47.7	
55-64	200	16,683	131	68.2	61.1-75.3	69	31.8	24.7-38.9	
65+	267	20,845	146	54.9	48.2-61.6	121	45.1	38.4-51.8	
Less Than H.S.	61	8,062	37	62.6	48.9-76.3	24	37.4	23.7-51.1	
H.S. or G.E.D.	200	20,038	120	62.7	53.7-71.7	80	37.3	28.3-46.3	
Some Post-H.S.	179	24,961	106	56.8	47.0-66.6	73	43.2	33.4-53.0	
College Graduate	194	13,879	127	68.4	60.8-76.0	67	31.6	24.0-39.2	
\$15,000- 24,999	69	7,479	43	56.7	40.4-73.0	26	43.3	27.0-59.6	
\$25,000- 34,999	81	9,384	45	53.6	37.1-70.1	36	46.4	29.9-62.9	
\$35,000- 49,999	114	10,573	75	64.5	54.3-74.7	39	35.5	25.3-45.7	
\$50,000-74,999	113	13,029	75	63.8	51.5-76.1	38	36.2	23.9-48.5	
\$75,000+	158	15,424	96	64.2	55.6-72.8	62	35.8	27.2-44.4	

^{&#}x27;% = Percentage, C.I.(95%) = Confidence Interval (at 95 percent probability level). Percentages are weighted to population characteristics.

Rows with less than 50 observations have been suppressed.

TABLE 15: HAS A DOCTOR OR OTHER HEALTH PROFESSIONAL EVER ADVISED YOU TO DO ANY OF THE FOLLOWING TO HELP LOWER OR CONTROL YOUR HIGH BLOOD PRESSURE? (EVER ADVISED YOU TO) CHANGE YOUR EATING HABITS (TO HELP LOWER OR CONTROL YOUR HIGH BLOOD PRESSURE)? (BPEATADV)

Denominator excludes: Respondents who are male or respondents with refused or with missing responses

		ab weighteed	ubii.		INDIVI.			
DEMOGRAPHIC	RESPONDE	NT NUMBER		Yes	i		No	
GROUPS	TOTAL	WEIGHTED	N	%	C.I. (95%)	N	%	C.I. (95%)
Female	955	61,793	524	57.5	53.8-61.2	431	42.5	38.8-46.2
Non-Native	926	59,240	505	56.6	52.9-60.3	421	43.4	39.7-47.1
45-54	112	9,890	75	68.8	59.2-78.4	37	31.2	21.6-40.8
55-64	234	14,021	157	66.6	59.9-73.3	77	33.4	26.7-40.1
65+	544	28,896	257	49.2	44.5-53.9	287	50.8	46.1-55.5
Less Than H.S.	76	6,562	31	42.8	30.3-55.3	45	57.2	44.7-69.7
H.S. or G.E.D.	330	19,731	182	57.8	51.3-64.3	148	42.2	35.7-48.7
Some Post-H.S.	299	23,005	171	61.5	55.4-67.6	128	38.5	32.4-44.6
College Graduate	248	12,413	140	57.6	49.8-65.4	108	42.4	34.6-50.2
Less than \$15,000	103	7,654	54	59.2	48.2-70.2	49	40.8	29.8-51.8
\$15,000- 24,999	184	12,230	99	57.8	49.2-66.4	85	42.2	33.6-50.8
\$25,000- 34,999	114	6,019	60	56.9	46.9-66.9	54	43.1	33.1-53.1
\$35,000- 49,999	118	7,251	59	50.5	38.9-62.1	59	49.5	37.9-61.1
\$50,000- 74,999	105	6,936	61	62.9	52.5-73.3	44	37.1	26.7-47.5
\$75,000+	120	8,283	74	64.0	53.6-74.4	46	36.0	25.6-46.4

NOTE:

^{&#}x27;% = Percentage, C.I.(95%) = Confidence Interval (at 95 percent probability level). Percentages are weighted to population characteristics.

Rows with less than 50 observations have been suppressed.

TABLE 16: [HAS A DOCTOR OR OTHER HEALTH PROFESSIONAL EVER ADVISED YOU TO DO ANY OF THE FOLLOWING TO HELP LOWER OR CONTROL YOUR HIGH BLOOD PRESSURE?] (EVER ADVISED YOU TO) CUT DOWN ON SALT (TO HELP LOWER OR CONTROL YOUR HIGH BLOOD PRESSURE)? (BPSLTADV)

Denominator excludes: Respondents with refused/missing responses

Data was weighted using LANDWT.

	Data W	was weighted using _LIANDWI.									
DEMOGRAPHIC	RESPONDE	RESPONDENT NUMBER		Yes			No		Do	not u	se salt
GROUPS	TOTAL	WEIGHTED	N	%	C.I. (95%)	N	%	C.I. (95%)	N	%	C.I. (95%)
TOTAL	1594	129,421	1061	67.3	64.4-70.2	498	30.8	27.9-33.7	35	1.9	01.3-02.5
Male	642	67,582	431	67.2	62.3-72.1	197	31.1	26.4-35.8	14	1.7	00.7-02.7
Female	952	61,839	630	67.5	64.0-71.0	301	30.5	27.0-34.0	21	2.0	01.0-03.0
Non-Native	1543	124,424	1024	66.8	63.7-69.9	486	31.4	28.3-34.5	33	1.8	01.2-02.4
35-44	77	12,867	47	58.9	45.4-72.4	30	41.1	27.6-54.6			
45-54	211	22,827	142	68.8	61.7-75.9	66	30.0	22.9-37.1	3	1.1	00.0-02.5
55-64	432	30,606	296	69.0	64.1-73.9	130	29.5	24.6-34.4	6	1.4	00.0-02.8
65+	817	50,576	540	67.7	64.2-71.2	251	28.9	25.6-32.2	26	3.4	02.0-04.8
Less Than H.S.	136	14,669	90	66.2	57.0-75.4	41	30.0	21.0-39.0	5	3.8	00.5-07.1
H.S. or G.E.D.	529	39,550	357	68.1	62.8-73.4	156	29.1	24.0-34.2	16	2.7	01.3-04.1
Some Post-H.S.	484	48,585	324	66.9	61.4-72.4	153	32.1	26.6-37.6	7	1.0	00.2-01.8
College Graduate	441	26,192	287	68.1	63.0-73.2	147	30.8	25.7-35.9	7	1.1	00.3-01.9
Less than \$15,000	142	12,172	92	66.4	57.2-75.6	44	29.8	20.8-38.8	6	3.8	00.3-07.3
\$15,000- 24,999	252	19,710	177	68.3	60.1-76.5	72	30.6	22.6-38.6	3	1.2	00.0-02.6
\$25,000- 34,999	194	15,368	136	75.1	67.7-82.5	51	22.5	15.4-29.6	7	2.4	00.6-04.2
\$35,000- 49,999	236	18,199	153	65.1	57.7-72.5	79	33.5	26.1-40.9	4	1.3	00.0-02.7
\$50,000- 74,999	219	20,037	142	62.7	53.7-71.7	74	36.5	27.5-45.5	3	8.0	00.0-01.8
\$75.000+	279	23.805	182	67.4	60.9-73.9	93	31.0	24.5-37.5	4	1.6	00.0-03.2

^{&#}x27;% = Percentage, C.I.(95%) = Confidence Interval (at 95 percent probability level). Percentages are weighted to population characteristics.

Rows with less than 50 observations have been suppressed.

TABLE 17: [HAS A DOCTOR OR OTHER HEALTH PROFESSIONAL EVER ADVISED YOU TO DO ANY OF THE FOLLOWING TO HELP LOWER OR CONTROL YOUR HIGH BLOOD PRESSURE?] (EVER ADVISED YOU TO) CUT DOWN ON SALT (TO HELP LOWER OR CONTROL YOUR HIGH BLOOD PRESSURE)? (BPSLTADV)

Denominator excludes: Respondents who are female or respondents with refused or with missing responses

Data was weighted using _LANDWT.

				J			-				
DEMOGRAPHIC	RESPONDENT NUMBER		Yes			No			Do not use salt		se salt
GROUPS	TOTAL	WEIGHTED	N	%	C.I. (95%)	N	%	C.I. (95%)	N	%	C.I. (95%)
Male	642	67,582	431	67.2	62.3-72.1	197	31.1	26.4-35.8	14	1.7	00.7-02.7
Non-Native	620	65,138	416	66.9	62.0-71.8	191	31.5	26.6-36.4	13	1.6	00.6-02.6
45-54	99	12,937	64	67.4	57.4-77.4	32	30.6	20.6-40.6	3	2.0	00.0-04.4
55-64	201	16,780	138	69.8	62.7-76.9	61	29.6	22.5-36.7	2	0.6	00.0-01.4
65+	272	21,378	182	66.8	60.7-72.9	81	29.4	23.5-35.3	9	3.8	01.3-06.3
Less Than H.S.	61	8,142	40	68.2	55.5-80.9	19	28.8	16.3-41.3	2	3.0	00.0-07.1
H.S. or G.E.D.	204	20,057	141	70.4	62.4-78.4	56	27.0	19.2-34.8	7	2.6	00.6-04.6
Some Post-H.S.	182	25,214	118	63.7	54.5-72.9	61	35.1	25.9-44.3	3	1.1	00.0-02.5
College Graduate	193	13,825	131	69.4	62.0-76.8	60	29.8	22.4-37.2	2	0.8	00.0-02.0
\$15,000- 24,999	70	7,564	48	61.0	44.7-77.3	20	36.4	19.9-52.9	2	2.6	00.0-06.3
\$25,000- 34,999	82	9,474	60	79.8	70.2-89.4	19	18.2	09.0-27.4	3	2.0	00.0-04.4
\$35,000- 49,999	115	10,671	78	68.7	59.3-78.1	33	29.0	19.6-38.4	4	2.3	00.0-04.7
\$50,000- 74,999	114	13,102	74	59.4	46.9-71.9	39	40.0	27.5-52.5	1	0.6	00.0-01.8
\$75,000+	159	15,521	104	67.4	59.0-75.8	52	30.5	22.3-38.7	3	2.1	00.0-04.5

^{&#}x27;% = Percentage, C.I.(95%) = Confidence Interval (at 95 percent probability level). Percentages are weighted to population characteristics.

Rows with less than 50 observations have been suppressed.

TABLE 18: [HAS A DOCTOR OR OTHER HEALTH PROFESSIONAL EVER ADVISED YOU TO DO ANY OF THE FOLLOWING TO HELP LOWER OR CONTROL YOUR HIGH BLOOD PRESSURE?] (EVER ADVISED YOU TO) CUT DOWN ON SALT (TO HELP LOWER OR CONTROL YOUR HIGH BLOOD PRESSURE)? (BPSLTADV)

Denominator excludes: Respondents who are male or respondents with refused or with missing responses

Data was weighted using _LANDWT.

DEMOGRAPHIC	RESPONDE	Yes			No	ı	Do not use salt				
GROUPS	TOTAL	WEIGHTED	N	%	C.I. (95%)	N	%	C.I. (95%)	N	%	C.I. (95%)
Female	952	61,839	630	67.5	64.0-71.0	301	30.5	27.0-34.0	21	2.0	01.0-03.0
Non-Native	923	59,286	608	66.6	62.9-70.3	295	31.3	27.8-34.8	20	2.1	01.1-03.1
45-54	112	9,890	78	70.8	61.2-80.4	34	29.2	19.6-38.8			
55-64	231	13,825	158	68.2	61.5-74.9	69	29.4	23.1-35.7	4	2.5	00.0-05.2
65+	545	29,198	358	68.3	64.2-72.4	170	28.6	24.5-32.7	17	3.1	01.5-04.7
Less Than H.S.	75	6,527	50	63.7	51.0-76.4	22	31.4	18.9-43.9	3	4.9	00.0-10.4
H.S. or G.E.D.	325	19,493	216	65.8	59.3-72.3	100	31.4	25.1-37.7	9	2.8	00.8-04.8
Some Post-H.S.	302	23,371	206	70.2	64.5-75.9	92	28.8	23.1-34.5	4	0.9	00.0-01.9
College Graduate	248	12,367	156	66.7	59.6-73.8	87	31.9	25.0-38.8	5	1.4	00.2-02.6
Less than \$15,000	102	7,568	65	68.1	57.9-78.3	31	25.8	16.4-35.2	6	6.1	00.6-11.6
\$15,000- 24,999	182	12,146	129	72.8	65.4-80.2	52	26.9	19.5-34.3	1	0.3	00.0-00.9
\$25,000- 34,999	112	5,894	76	67.5	57.7-77.3	32	29.4	19.8-39.0	4	3.1	00.0-06.2
\$35,000- 49,999	121	7,529	75	60.0	48.2-71.8	46	40.0	28.2-51.8			
\$50,000- 74,999	105	6,936	68	68.8	59.0-78.6	35	29.9	20.1-39.7	2	1.3	00.0-03.5
\$75,000+	120	8,283	78	67.3	57.5-77.1	41	32.0	22.2-41.8	1	0.7	00.0-02.1

^{&#}x27;% = Percentage, C.I.(95%) = Confidence Interval (at 95 percent probability level). Percentages are weighted to population characteristics.

Rows with less than 50 observations have been suppressed.

TABLE 19: [HAS A DOCTOR OR OTHER HEALTH PROFESSIONAL EVER ADVISED YOU TO DO ANY OF THE FOLLOWING TO HELP LOWER OR CONTROL YOUR HIGH BLOOD PRESSURE?] (EVER ADVISED YOU TO) REDUCE ALCOHOL USE (TO HELP LOWER OR CONTROL YOUR HIGH BLOOD PRESSURE)? (BPALCADV)

Denominator excludes: Respondents with refused/missing responses

Data was weighted using _LANDWT.

		Data wa	ab WCI	gircea	using _r	. דוא מאדאו	•				
DEMOGRAPHIC	RESPONDE	NT NUMBER	Yes				No			Do not	drink
GROUPS	TOTAL	WEIGHTED	N	%	C.I. (95%)	N	%	C.I. (95%)	N	%	C.I. (95%)
TOTAL	1600	129,982	396	28.1	25.0-31.2	915	55.5	52.2-58.8	289	16.4	14.0-18.8
Male	644	68,020	210	34.4	29.3-39.5	347	53.1	47.8-58.4	87	12.4	09.1-15.7
Female	956	61,962	186	21.1	18.0-24.2	568	58.2	54.5-61.9	202	20.8	17.9-23.7
Non-Native	1549	124,985	379	27.7	24.6-30.8	896	56.3	53.0-59.6	274	15.9	13.5-18.3
35-44	77	12,867	30	41.0	27.3-54.7	42	47.9	34.4-61.4	5	11.0	00.0-22.2
45-54	212	23,225	70	31.5	24.6-38.4	119	57.3	49.7-64.9	23	11.2	06.1-16.3
55-64	435	30,791	133	32.5	27.4-37.6	247	54.8	49.5-60.1	55	12.7	09.2-16.2
65+	819	50,578	147	19.4	16.3-22.5	476	56.6	52.7-60.5	196	24.0	20.7-27.3
Less Than H.S.	139	14,867	27	26.1	16.5-35.7	70	47.9	38.3-57.5	42	26.0	18.2-33.8
H.S. or G.E.D.	531	39,902	124	24.2	19.7-28.7	295	58.2	52.7-63.7	112	17.7	14.2-21.2
Some Post-H.S.	484	48,530	140	32.1	26.4-37.8	275	53.7	47.8-59.6	69	14.2	09.9-18.5
College Graduate	442	26,258	105	28.1	22.2-34.0	274	59.3	53.2-65.4	63	12.7	09.0-16.4
Less than \$15,000	143	12,258	38	29.1	20.3-37.9	74	51.2	41.4-61.0	31	19.7	12.1-27.3
\$15,000- 24,999	252	19,663	68	29.2	21.9-36.5	126	49.5	41.5-57.5	58	21.3	15.0-27.6
\$25,000- 34,999	196	15,435	47	31.7	19.9-43.5	110	54.4	43.4-65.4	39	13.9	08.6-19.2
\$35,000- 49,999	236	18,199	55	24.2	17.7-30.7	143	58.0	50.4-65.6	38	17.8	11.3-24.3
\$50,000- 74,999	218	19,962	69	34.8	26.2-43.4	124	52.8	43.8-61.8	25	12.4	05.1-19.7
\$75.000+	281	23.900	72	31.0	23.9-38.1	180	60.1	52.8-67.4	29	9.0	05.5-12.5

^{&#}x27;% = Percentage, C.I.(95%) = Confidence Interval (at 95 percent probability level). Percentages are weighted to population characteristics.

Rows with less than 50 observations have been suppressed.

TABLE 20: [HAS A DOCTOR OR OTHER HEALTH PROFESSIONAL EVER ADVISED YOU TO DO ANY OF THE FOLLOWING TO HELP LOWER OR CONTROL YOUR HIGH BLOOD PRESSURE?] (EVER ADVISED YOU TO) REDUCE ALCOHOL USE (TO HELP LOWER OR CONTROL YOUR HIGH BLOOD PRESSURE)? (BPALCADV)

Denominator excludes: Respondents who are female or respondents with refused or with missing responses

Data was weighted using _LANDWT.

DEMOGRAPHIC	RESPONDENT NUMBER		Yes			No			Do not drink		
GROUPS	TOTAL	WEIGHTED	N	%	C.I. (95%)	N	%	C.I. (95%)	N	%	C.I. (95%)
Male	644	68,020	210	34.4	29.3-39.5	347	53.1	47.8-58.4	87	12.4	09.1-15.7
Non-Native	622	65,576	197	33.4	28.3-38.5	341	54.3	49.0-59.6	84	12.3	09.0-15.6
45-54	100	13,335	37	32.0	22.2-41.8	53	56.6	45.6-67.6	10	11.4	04.0-18.8
55-64	201	16,759	76	38.4	31.0-45.8	101	50.2	42.6-57.8	24	11.4	06.5-16.3
65+	273	21,440	68	26.5	20.6-32.4	157	56.4	49.9-62.9	48	17.1	12.2-22.0
Less Than H.S.	62	8,203	17	33.6	19.1-48.1	35	53.4	38.9-67.9	10	13.0	04.4-21.6
H.S. or G.E.D.	204	20,380	64	29.5	21.9-37.1	106	56.7	47.7-65.7	34	13.8	08.7-18.9
Some Post-H.S.	182	25,214	68	38.2	28.8-47.6	93	48.6	39.2-58.0	21	13.2	06.3-20.1
College Graduate	194	13,879	61	36.2	27.0-45.4	112	55.6	46.6-64.6	21	8.3	04.6-12.0
\$15,000- 24,999	70	7,564	30	38.9	24.6-53.2	26	41.9	25.8-58.0	14	19.2	08.4-30.0
\$25,000- 34,999	82	9,474	28	41.7	24.8-58.6	41	48.1	32.2-64.0	13	10.2	03.5-16.9
\$35,000- 49,999	115	10,671	35	29.8	20.2-39.4	62	51.8	41.2-62.4	18	18.3	09.1-27.5
\$50,000- 74,999	113	13,026	42	38.5	26.5-50.5	57	48.5	36.2-60.8	14	13.0	02.6-23.4
\$75.000+	160	15.575	48	35.1	25.5-44.7	98	57.6	48.0-67.2	14	7.3	03.2-11.4

^{&#}x27;% = Percentage, C.I.(95%) = Confidence Interval (at 95 percent probability level). Percentages are weighted to population characteristics.

Rows with less than 50 observations have been suppressed.

TABLE 21: [HAS A DOCTOR OR OTHER HEALTH PROFESSIONAL EVER ADVISED YOU TO DO ANY OF THE FOLLOWING TO HELP LOWER OR CONTROL YOUR HIGH BLOOD PRESSURE?] (EVER ADVISED YOU TO) REDUCE ALCOHOL USE (TO HELP LOWER OR CONTROL YOUR HIGH BLOOD PRESSURE)? (BPALCADV)

Denominator excludes: Respondents who are male or respondents with refused or with missing responses

Data was weighted using _LANDWT.

DEMOGRAPHIC	RESPONDE	Yes			No			Do not drink			
GROUPS	TOTAL	WEIGHTED	N	%	C.I. (95%)	N	%	C.I. (95%)	N	%	C.I. (95%)
Female	956	61,962	186	21.1	18.0-24.2	568	58.2	54.5-61.9	202	20.8	17.9-23.7
Non-Native	927	59,409	182	21.5	18.2-24.8	555	58.6	54.9-62.3	190	19.9	17.0-22.8
45-54	112	9,890	33	30.8	21.4-40.2	66	58.2	48.0-68.4	13	11.0	04.5-17.5
55-64	234	14,033	57	25.6	19.3-31.9	146	60.2	53.1-67.3	31	14.2	09.1-19.3
65+	546	29,139	79	14.2	11.1-17.3	319	56.7	52.2-61.2	148	29.1	24.8-33.4
Less Than H.S.	77	6,663	10	16.8	05.6-28.0	35	41.2	29.0-53.4	32	42.0	29.7-54.3
H.S. or G.E.D.	327	19,522	60	18.6	13.7-23.5	189	59.7	53.4-66.0	78	21.7	17.0-26.4
Some Post-H.S.	302	23,315	72	25.5	19.8-31.2	182	59.2	52.7-65.7	48	15.3	10.6-20.0
College Graduate	248	12,379	44	19.0	12.7-25.3	162	63.4	55.6-71.2	42	17.6	10.9-24.3
Less than \$15,000	103	7,654	23	24.0	14.6-33.4	53	48.3	36.7-59.9	27	27.7	16.9-38.5
\$15,000- 24,999	182	12,099	38	23.1	15.1-31.1	100	54.3	45.3-63.3	44	22.6	15.2-30.0
\$25,000- 34,999	114	5,961	19	15.9	08.6-23.2	69	64.4	54.8-74.0	26	19.7	12.1-27.3
\$35,000- 49,999	121	7,529	20	16.1	08.8-23.4	81	66.7	56.5-76.9	20	17.2	09.0-25.4
\$50,000-74,999	105	6,936	27	27.8	17.2-38.4	67	61.0	49.8-72.2	11	11.2	04.1-18.3
\$75,000+	121	8,324	24	23.2	14.2-32.2	82	64.7	54.7-74.7	15	12.0	05.1-18.9

^{&#}x27;% = Percentage, C.I.(95%) = Confidence Interval (at 95 percent probability level). Percentages are weighted to population characteristics.

Rows with less than 50 observations have been suppressed.

TABLE 22: [HAS A DOCTOR OR OTHER HEALTH PROFESSIONAL EVER ADVISED YOU TO DO ANY OF THE FOLLOWING TO HELP LOWER OR CONTROL YOUR HIGH BLOOD PRESSURE?] (EVER ADVISED YOU TO) EXERCISE (TO HELP LOWER OR CONTROL YOUR HIGH BLOOD PRESSURE)? (BPEXRADV)

Denominator excludes: Respondents with refused/missing responses

Data was weighted using _LANDWT.

DEMOGRAPHIC	RESPONDE	NT NUMBER		Yes	·· - ·	No			
GROUPS	TOTAL	WEIGHTED	N	%	C.I. (95%)	N	%	C.I. (95%)	
TOTAL	1591	129,169	1153	74.0	71.1-76.9	438	26.0	23.1-28.9	
Male	639	67,527	479	74.5	69.6-79.4	160	25.5	20.6-30.4	
Female	952	61,642	674	73.5	70.4-76.6	278	26.5	23.4-29.6	
Non-Native	1542	124,354	1114	73.5	70.4-76.6	428	26.5	23.4-29.6	
35-44	77	12,867	60	80.1	70.1-90.1	17	19.9	09.9-29.9	
45-54	212	23,225	161	76.3	69.8-82.8	51	23.7	17.2-30.2	
55-64	434	30,713	340	78.0	73.5-82.5	94	22.0	17.5-26.5	
65+	811	49,920	553	70.2	66.7-73.7	258	29.8	26.3-33.3	
Less Than H.S.	137	14,602	93	71.3	62.9-79.7	44	28.7	20.3-37.1	
H.S. or G.E.D.	530	39,750	368	70.7	64.4-77.0	162	29.3	23.0-35.6	
Some Post-H.S.	482	48,320	349	75.5	70.8-80.2	133	24.5	19.8-29.2	
College Graduate	438	26,072	342	78.8	74.3-83.3	96	21.2	16.7-25.7	
Less than \$15,000	142	12,172	97	71.2	62.4-80.0	45	28.8	20.0-37.6	
\$15,000- 24,999	251	19,423	175	68.0	59.8-76.2	76	32.0	23.8-40.2	
\$25,000- 34,999	193	15,267	137	78.4	71.7-85.1	56	21.6	14.9-28.3	
\$35,000- 49,999	234	18,101	166	71.3	64.4-78.2	68	28.7	21.8-35.6	
\$50,000- 74,999	219	20,037	174	82.4	76.1-88.7	45	17.6	11.3-23.9	
\$75,000+	281	23,900	213	77.9	72.6-83.2	68	22.1	16.8-27.4	

^{&#}x27;% = Percentage, C.I.(95%) = Confidence Interval (at 95 percent probability level). Percentages are weighted to population characteristics.

Rows with less than 50 observations have been suppressed.

TABLE 23: [HAS A DOCTOR OR OTHER HEALTH PROFESSIONAL EVER ADVISED YOU TO DO ANY OF THE FOLLOWING TO HELP LOWER OR CONTROL YOUR HIGH BLOOD PRESSURE?] (EVER ADVISED YOU TO) EXERCISE (TO HELP LOWER OR CONTROL YOUR HIGH BLOOD PRESSURE)? (BPEXRADV)

Denominator excludes: Respondents who are female or respondents with refused or with missing responses

Data was weighted using _LANDWT.

		-		·					
DEMOGRAPHIC	RESPONDE	NT NUMBER		Yes	6	No			
GROUPS	TOTAL	WEIGHTED	N	%	C.I. (95%)	N	%	C.I. (95%)	
Male	639	67,527	479	74.5	69.6-79.4	160	25.5	20.6-30.4	
Non-Native	618	65,245	461	73.9	68.8-79.0	157	26.1	21.0-31.2	
45-54	100	13,335	74	76.8	68.2-85.4	26	23.2	14.6-31.8	
55-64	201	16,778	155	76.7	70.0-83.4	46	23.3	16.6-30.0	
65+	269	21,090	197	73.3	67.6-79.0	72	26.7	21.0-32.4	
Less Than H.S.	62	8,203	44	73.9	61.9-85.9	18	26.1	14.1-38.1	
H.S. or G.E.D.	202	20,172	145	69.0	58.2-79.8	57	31.0	20.2-41.8	
Some Post-H.S.	181	25,042	132	75.8	68.0-83.6	49	24.2	16.4-32.0	
College Graduate	192	13,766	158	82.3	76.0-88.6	34	17.7	11.4-24.0	
\$15,000- 24,999	69	7,392	50	63.2	46.3-80.1	19	36.8	19.9-53.7	
\$25,000- 34,999	82	9,474	61	82.9	74.5-91.3	21	17.1	08.7-25.5	
\$35,000- 49,999	113	10,572	81	69.8	60.2-79.4	32	30.2	20.6-39.8	
\$50,000- 74,999	114	13,102	93	83.3	74.9-91.7	21	16.7	08.3-25.1	
\$75,000+	160	15,575	119	77.1	70.0-84.2	41	22.9	15.8-30.0	

^{&#}x27;% = Percentage, C.I.(95%) = Confidence Interval (at 95 percent probability level). Percentages are weighted to population characteristics.

Rows with less than 50 observations have been suppressed.

TABLE 24: [HAS A DOCTOR OR OTHER HEALTH PROFESSIONAL EVER ADVISED YOU TO DO ANY OF THE FOLLOWING TO HELP LOWER OR CONTROL YOUR HIGH BLOOD PRESSURE?] (EVER ADVISED YOU TO) EXERCISE (TO HELP LOWER OR CONTROL YOUR HIGH BLOOD PRESSURE)? (BPEXRADV)

Denominator excludes: Respondents who are male or respondents with refused or with missing responses

Data was weighted using _LANDWT.

DEMOGRAPHIC	RESPONDE	NT NUMBER	u uzii	Yes	3	No			
GROUPS	TOTAL	WEIGHTED	N	%	C.I. (95%)	N	%	C.I. (95%)	
Female	952	61,642	674	73.5	70.4-76.6	278	26.5	23.4-29.6	
Non-Native	924	59,109	653	73.0	69.7-76.3	271	27.0	23.7-30.3	
45-54	112	9,890	87	75.6	66.2-85.0	25	24.4	15.0-33.8	
55-64	233	13,935	185	79.5	73.8-85.2	48	20.5	14.8-26.2	
65+	542	28,831	356	68.0	63.9-72.1	186	32.0	27.9-36.1	
Less Than H.S.	75	6,399	49	67.9	56.1-79.7	26	32.1	20.3-43.9	
H.S. or G.E.D.	328	19,577	223	72.4	67.1-77.7	105	27.6	22.3-32.9	
Some Post-H.S.	301	23,278	217	75.2	69.9-80.5	84	24.8	19.5-30.1	
College Graduate	246	12,306	184	75.0	68.3-81.7	62	25.0	18.3-31.7	
Less than \$15,000	102	7,568	66	69.0	58.8-79.2	36	31.0	20.8-41.2	
\$15,000- 24,999	182	12,032	125	71.0	63.2-78.8	57	29.0	21.2-36.8	
\$25,000- 34,999	111	5,793	76	71.0	62.0-80.0	35	29.0	20.0-38.0	
\$35,000- 49,999	121	7,529	85	73.4	64.2-82.6	36	26.6	17.4-35.8	
\$50,000- 74,999	105	6,936	81	80.7	72.7-88.7	24	19.3	11.3-27.3	
\$75,000+	121	8,324	94	79.4	71.6-87.2	27	20.6	12.8-28.4	

^{&#}x27;% = Percentage, C.I.(95%) = Confidence Interval (at 95 percent probability level). Percentages are weighted to population characteristics.

Rows with less than 50 observations have been suppressed.

TABLE 25: [HAS A DOCTOR OR OTHER HEALTH PROFESSIONAL EVER ADVISED YOU TO DO ANY OF THE FOLLOWING TO HELP LOWER OR CONTROL YOUR HIGH BLOOD PRESSURE?] (EVER ADVISED YOU TO) TAKE MEDICATION (TO HELP LOWER OR CONTROL YOUR HIGH BLOOD PRESSURE)? (BPMEDADV)

Denominator excludes: Respondents with refused/missing responses

Data was weighted using _LANDWT.

	Data W	as weighte	u usiii	.g	TADAAT.				
DEMOGRAPHIC	RESPONDE	NT NUMBER		Yes	3		No)	
GROUPS	TOTAL	WEIGHTED	N	%	C.I. (95%)	N	%	C.I. (95%)	
TOTAL	1604	130,243	1472	87.9	85.0-90.8	132	12.1	09.2-15.0	
Male	644	68,033	577	84.6	79.7-89.5	67	15.4	10.5-20.3	
Female	960	62,210	895	91.5	89.0-94.0	65	8.5	06.0-11.0	
Non-Native	1554	125,308	1426	87.8	84.9-90.7	128	12.2	09.3-15.1	
35-44	77	12,867	57	75.1	64.1-86.1	20	24.9	13.9-35.9	
45-54	210	23,054	182	87.9	83.2-92.6	28	12.1	07.4-16.8	
55-64	437	30,896	403	91.7	88.8-94.6	34	8.3	05.4-11.2	
65+	822	50,820	788	95.9	94.3-97.5	34	4.1	02.5-05.7	
Less Than H.S.	139	14,867	134	96.6	93.5-99.7	5	3.4	00.3-06.5	
H.S. or G.E.D.	531	39,886	488	84.9	78.2-91.6	43	15.1	08.4-21.8	
Some Post-H.S.	486	48,693	438	86.7	82.0-91.4	48	13.3	08.6-18.0	
College Graduate	444	26,372	408	89.5	85.6-93.4	36	10.5	06.6-14.4	
Less than \$15,000	143	12,258	127	88.9	83.0-94.8	16	11.1	05.2-17.0	
\$15,000- 24,999	254	19,795	234	89.6	84.3-94.9	20	10.4	05.1-15.7	
\$25,000- 34,999	196	15,435	181	87.1	76.5-97.7	15	12.9	02.3-23.5	
\$35,000-49,999	236	18,199	226	92.6	87.3-97.9	10	7.4	02.1-12.7	
\$50,000-74,999	219	20,037	192	81.8	74.2-89.4	27	18.2	10.6-25.8	
\$75,000+	280	23,792	253	88.5	84.0-93.0	27	11.5	07.0-16.0	

^{&#}x27;% = Percentage, C.I.(95%) = Confidence Interval (at 95 percent probability level). Percentages are weighted to population characteristics.

Rows with less than 50 observations have been suppressed.

TABLE 26: [HAS A DOCTOR OR OTHER HEALTH PROFESSIONAL EVER ADVISED YOU TO DO ANY OF THE FOLLOWING TO HELP LOWER OR CONTROL YOUR HIGH BLOOD PRESSURE?] (EVER ADVISED YOU TO) TAKE MEDICATION (TO HELP LOWER OR CONTROL YOUR HIGH BLOOD PRESSURE)? (BPMEDADV)

Denominator excludes: Respondents who are female or respondents with refused or with missing responses

Data was weighted using _LANDWT.

		_						
DEMOGRAPHIC	RESPONDENT NUMBER			Yes	3		No	
GROUPS	TOTAL	WEIGHTED	N	%	C.I. (95%)	N	%	C.I. (95%)
Male	644	68,033	577	84.6	79.7-89.5	67	15.4	10.5-20.3
Non-Native	623	65,651	557	84.2	79.1-89.3	66	15.8	10.7-20.9
45-54	99	13,272	86	88.6	82.3-94.9	13	11.4	05.1-17.7
55-64	202	16,834	185	90.5	85.8-95.2	17	9.5	04.8-14.2
65+	273	21,440	259	94.9	92.2-97.6	14	5.1	02.4-07.8
Less Than H.S.	62	8,203	59	95.5	90.4-0100	3	4.5	00.0-09.6
H.S. or G.E.D.	204	20,393	182	79.0	67.2-90.8	22	21.0	09.2-32.8
Some Post-H.S.	182	25,214	156	83.0	75.0-91.0	26	17.0	09.0-25.0
College Graduate	194	13,879	178	88.8	82.9-94.7	16	11.2	05.3-17.1
\$15,000- 24,999	70	7,564	61	84.9	74.3-95.5	9	15.1	04.5-25.7
\$25,000- 34,999	82	9,474	74	83.0	66.5-99.5	8	17.0	00.5-33.5
\$35,000- 49,999	115	10,671	110	94.1	88.8-99.4	5	5.9	00.6-11.2
\$50,000-74,999	114	13,102	97	77.1	66.1-88.1	17	22.9	11.9-33.9
\$75,000+	160	15,575	144	88.9	83.4-94.4	16	11.1	05.6-16.6

^{&#}x27;% = Percentage, C.I.(95%) = Confidence Interval (at 95 percent probability level). Percentages are weighted to population characteristics.

Rows with less than 50 observations have been suppressed.

TABLE 27: [HAS A DOCTOR OR OTHER HEALTH PROFESSIONAL EVER ADVISED YOU TO DO ANY OF THE FOLLOWING TO HELP LOWER OR CONTROL YOUR HIGH BLOOD PRESSURE?] (EVER ADVISED YOU TO) TAKE MEDICATION (TO HELP LOWER OR CONTROL YOUR HIGH BLOOD PRESSURE)? (BPMEDADV)

Denominator excludes: Respondents who are male or respondents with refused or with missing responses

DEMOGRAPHIC	RESPONDE	NT NUMBER	a abiii	Yes	3	No				
GROUPS	TOTAL	WEIGHTED	N	%	C.I. (95%)	N	%	C.I. (95%)		
Female	960	62,210	895	91.5	89.0-94.0	65	8.5	06.0-11.0		
Non-Native	931	59,657	869	91.9	89.5-94.3	62	8.1	05.7-10.5		
45-54	111	9,782	96	87.0	79.7-94.3	15	13.0	05.7-20.3		
55-64	235	14,062	218	93.1	89.8-96.4	17	6.9	03.6-10.2		
65+	549	29,380	529	96.6	95.0-98.2	20	3.4	01.8-05.0		
Less Than H.S.	77	6,663	75	97.9	95.0-0100	2	2.1	00.0-05.0		
H.S. or G.E.D.	327	19,492	306	91.1	86.2-96.0	21	8.9	04.0-13.8		
Some Post-H.S.	304	23,479	282	90.7	86.4-95.0	22	9.3	05.0-13.6		
College Graduate	250	12,494	230	90.3	85.2-95.4	20	9.7	04.6-14.8		
Less than \$15,000	103	7,654	93	91.0	84.9-97.1	10	9.0	02.9-15.1		
\$15,000- 24,999	184	12,230	173	92.5	86.8-98.2	11	7.5	01.8-13.2		
\$25,000- 34,999	114	5,961	107	93.7	88.8-98.6	7	6.3	01.4-11.2		
\$35,000- 49,999	121	7,529	116	90.6	80.4-0100	5	9.4	00.0-19.6		
\$50,000- 74,999	105	6,936	95	90.6	84.1-97.1	10	9.4	02.9-15.9		
\$75,000+	120	8,216	109	87.7	79.9-95.5	11	12.3	04.5-20.1		

NOTE:

^{&#}x27;% = Percentage, C.I.(95%) = Confidence Interval (at 95 percent probability level). Percentages are weighted to population characteristics.

Rows with less than 50 observations have been suppressed.

TABLE 28: WERE YOU TOLD ON TWO OR MORE DIFFERENT VISITS TO A DOCTOR OR OTHER HEALTH PROFESSIONAL THAT YOU HAD HIGH BLOOD PRESSURE? (BPHI2MR) Denominator excludes: Respondents with refused/missing responses

Data was weighted using _LANDWT.

DEMOGRAPHIC	RESPONDE	NT NUMBER	Yes		Yes, during pregnancy			No					tensive	
GROUPS	TOTAL	WEIGHTED	N	%	C.I. (95%)	N	%	C.I. (95%)	N	%	C.I. (95%)	N	%	C.I. (95%)
TOTAL	1574	128,337	1256	78.6	75.5-81.7	1	0.0	00.0-00.0	307	20.8	17.9-23.7	10	0.7	00.3-01.1
Male	634	66,975	508	77.6	72.5-82.7				120	21.4	16.3-26.5	6	1.0	00.2-01.8
Female	940	61,362	748	79.6	76.5-82.7	1	0.1	00.0-00.3	187	20.1	17.0-23.2	4	0.3	00.0-00.7
Non-Native	1525	123,403	1215	78.4	75.3-81.5				300	21.0	17.9-24.1	10	0.7	00.1-01.3
35-44	77	12,867	55	73.5	62.3-84.7				22	26.5	15.3-37.7			
45-54	209	22,718	157	76.1	69.6-82.6				49	22.1	15.8-28.4	3	1.8	00.0-04.0
55-64	428	30,425	347	82.2	78.3-86.1				79	17.3	13.4-21.2	2	0.6	00.0-01.4
65+	802	49,722	655	82.1	79.2-85.0	1	0.1	00.0-00.3	141	17.3	14.4-20.2	5	0.5	00.1-00.9
Less Than H.S.	137	14,683	114	84.8	78.1-91.5				23	15.2	08.5-21.9			
H.S. or G.E.D.	523	39,103	415	74.8	68.1-81.5				104	24.5	17.8-31.2	4	0.7	00.0-01.5
Some Post-H.S.	479	48,257	390	80.2	75.3-85.1				86	19.0	14.1-23.9	3	8.0	00.0-01.8
College Graduate	431	25,868	333	77.3	72.6-82.0	1	0.2	00.0-00.6	94	22.0	17.3-26.7	3	0.6	00.0-01.2
Less than \$15,000	141	12,177	118	85.6	79.3-91.9				23	14.4	08.1-20.7			
\$15,000- 24,999	249	19,473	198	81.9	76.6-87.2	1	0.2	00.0-00.6	47	16.6	11.5-21.7	3	1.4	00.0-03.2
\$25,000- 34,999	193	15,368	161	81.3	70.9-91.7				30	18.1	07.7-28.5	2	0.6	00.0-01.4
\$35,000- 49,999	234	17,969	194	83.3	77.2-89.4				39	16.4	10.3-22.5	1	0.3	00.0-00.9
\$50,000-74,999	217	19,925	168	75.1	67.1-83.1				49	24.9	16.9-32.9			
\$75,000+	274	23,544	206	74.0	67.7-80.3				65	24.6	18.5-30.7	3	1.3	00.0-03.1

NOTE:

Told Borderline

^{&#}x27;% = Percentage, C.I.(95%) = Confidence Interval (at 95 percent probability level). Percentages are weighted to population characteristics.

Rows with less than 50 observations have been suppressed.

TABLE 29: WERE YOU TOLD ON TWO OR MORE DIFFERENT VISITS TO A DOCTOR OR OTHER HEALTH PROFESSIONAL THAT YOU HAD HIGH BLOOD PRESSURE? (BPHI2MR) Denominator excludes: Respondents who are female or respondents with refused or with missing responses

Data was weighted using _LANDWT.

DEMOGRAPHIC	RESPONDENT NUMBER			Yes			No			old Bord re-hypert	,
GROUPS	TOTAL	WEIGHTED	N	%	C.I. (95%)	N	%	C.I. (95%)	N	%	C.I. (95%)
Male	634	66,975	508	77.6	72.5-82.7	120	21.4	16.3-26.5	6	1.0	00.2-01.8
Non-Native	612	64,531	489	77.5	72.2-82.8	117	21.5	16.4-26.6	6	1.0	00.0-02.0
45-54	99	12,937	73	74.1	64.5-83.7	24	23.1	13.9-32.3	2	2.8	00.0-06.7
55-64	199	16,657	167	85.7	80.6-90.8	30	13.3	08.2-18.4	2	1.0	00.0-02.6
65+	266	20,895	221	83.7	79.0-88.4	43	15.8	11.1-20.5	2	0.6	00.0-01.4
Less Than H.S.	60	8,020	53	90.6	83.3-97.9	7	9.4	02.1-16.7			
H.S. or G.E.D.	201	19,827	156	70.8	59.4-82.2	42	28.1	16.7-39.5	3	1.2	00.0-02.6
Some Post-H.S.	180	25,064	142	77.7	69.3-86.1	36	20.9	12.7-29.1	2	1.4	00.0-03.4
College Graduate	191	13,721	155	79.4	72.3-86.5	35	20.2	13.1-27.3	1	0.4	00.0-01.2
\$15,000- 24,999	69	7,467	50	77.7	66.9-88.5	17	19.5	09.5-29.5	2	2.8	00.0-07.3
\$25,000- 34,999	83	9,553	71	81.4	65.3-97.5	12	18.6	02.5-34.7			
\$35,000-49,999	113	10,440	99	87.0	79.9-94.1	13	12.5	05.4-19.6	1	0.5	00.0-01.5
\$50,000-74,999	113	13,026	85	70.3	58.9-81.7	28	29.7	18.3-41.1			
\$75,000+	157	15,412	126	77.2	69.4-85.0	29	21.1	13.7-28.5	2	1.7	00.0-04.2

^{&#}x27;% = Percentage, C.I.(95%) = Confidence Interval (at 95 percent probability level). Percentages are weighted to population characteristics.

Rows with less than 50 observations have been suppressed.

TABLE 30: WERE YOU TOLD ON TWO OR MORE DIFFERENT VISITS TO A DOCTOR OR OTHER HEALTH PROFESSIONAL THAT YOU HAD HIGH BLOOD PRESSURE? (BPHI2MR) Denominator excludes: Respondents who are male or respondents with refused or with missing responses

Data was weighted using _LANDWT.

DEMOGRAPHIC	RESPONDE	NT NUMBER	3, 3, 3,					No	1			derline, tensive		
GROUPS	TOTAL	WEIGHTED	N	%	C.I. (95%)	N	%	C.I. (95%)	N	%	C.I. (95%)	N	%	C.I. (95%)
Female	940	61,362	748	79.6	76.5-82.7	1	0.1	00.0-00.3	187	20.1	17.0-23.2	4	0.3	00.0-00.7
Non-Native	913	58,872	726	79.3	76.2-82.4				183	20.3	17.2-23.4	4	0.3	00.0-00.7
45-54	110	9,781	84	78.8	70.6-87.0				25	20.7	12.7-28.7	1	0.5	00.0-01.5
55-64	229	13,768	180	77.9	72.0-83.8				49	22.1	16.2-28.0			
65+	536	28,827	434	80.9	77.4-84.4	1	0.1	00.0-00.3	98	18.4	14.9-21.9	3	0.5	00.0-01.1
Less Than H.S.	77	6,663	61	77.8	66.4-89.2				16	22.2	10.8-33.6			
H.S. or G.E.D.	322	19,276	259	78.9	73.2-84.6				62	20.8	15.1-26.5	1	0.3	00.0-00.9
Some Post-H.S.	299	23,194	248	82.9	78.2-87.6				50	16.9	12.2-21.6	1	0.2	00.0-00.6
College Graduate	240	12,147	178	75.0	68.5-81.5	1	0.3	00.0-00.9	59	23.9	17.6-30.2	2	0.8	00.0-01.8
Less than \$15,000	101	7,573	86	86.1	78.7-93.5				15	13.9	06.5-21.3			
\$15,000- 24,999	180	12,006	148	84.4	78.9-89.9	1	0.3	00.0-00.9	30	14.8	09.5-20.1	1	0.5	00.0-01.5
\$25,000- 34,999	110	5,815	90	81.1	72.5-89.7				18	17.3	08.9-25.7	2	1.6	00.0-03.8
\$35,000- 49,999	121	7,529	95	78.1	67.7-88.5				26	21.9	11.5-32.3			
\$50,000-74,999	104	6,899	83	84.0	76.9-91.1				21	16.0	08.9-23.1			
\$75,000+	117	8,132	80	68.1	58.1-78.1				36	31.4	21.4-41.4	1	0.5	00.0-01.5

^{&#}x27;% = Percentage, C.I.(95%) = Confidence Interval (at 95 percent probability level). Percentages are weighted to population characteristics.

Rows with less than 50 observations have been suppressed.

BEHAVIORAL RISK FACTOR SURVEILLANCE SYSTEM NORTH DAKOTA MODULE QUESTIONS DATA REPORT, 2011

MODULE 32: RANDOM CHILD SELECTION CONTENTS

	HEN THE DENOMINATOR IS NOT BASED ON THE TOTAL POPULATION SURVEYED, THE TITLE CONTAINS A DESCRIPTION OF V	
IS	S EXCLUDED FROM THE DENOMINATOR. IF THE TITLE DOES NOT DESCRIBE THE DENOMINATOR, THE TABLE IS BASED ON A	LL RESPONDENTS
* * *	TO THE STATES SURVEY.	*****
TABLE		PAGE
111000	WHAT IS THE BIRTH MONTH AND YEAR OF THE "XTH" CHILD? (RCSBIRTH)	17101
1	MALES AND FEMALES	67
2	MALES ONLY	68
3	FEMALES ONLY	69
J	IS THE CHILD A BOY OR A GIRL? (RCSGENDR)	
4	MALES AND FEMALES	70
5	MALES ONLY	71
6	FEMALES ONLY	72
	IS THE CHILD HISPANIC OR LATINO? (RCHISLAT)	
7	MALES AND FEMALES	73
8	MALES ONLY	74
9	FEMALES ONLY	75
	WHICH ONE OR MORE OF THE FOLLOWING WOULD YOU SAY IS THE RACE OF THE CHILD? (RCSRACE)	
10	MALES AND FEMALES	76
11	MALES ONLY	77
12	FEMALES ONLY	78
	WHICH ONE OF THESE GROUPS WOULD YOU SAY BEST REPRESENTS THE CHILD'S RACE? (RCSBRACE)	
13	MALES AND FEMALES	79
14	MALES ONLY	80
15	FEMALES ONLY	81
	HOW ARE YOU RELATED TO THE CHILD? (RCSRLTN2)	
16	MALES AND FEMALES	82
17	MALES ONLY	83
18	FEMALES ONLY	84

67 of 93 July 26, 2012

TABLE 1: WHAT IS THE BIRTH MONTH AND YEAR OF THE "XTH" CHILD? (RCSBIRTH)

Denominator excludes: Respondents with do not know/refused/missing responses

Note: The tables for RCSBIRTH were produced using $_$ CHILDWT as the weighting variable

Denominator excludes respondents not reporting children

Data was weighted using _LANDWT.

DEMOGRAPHIC	RESPONDE	NT NUMBER		1993-1	997		1998-2	002		2003-2	007		2008-2	009		2010-2	011
GROUPS	TOTAL	WEIGHTED	N	%	C.I. (95%)	N	%	C.I. (95%)	N	%	C.I. (95%)	N	%	C.I. (95%)	N	%	C.I. (95%)
TOTAL	899	153,522	319	32.4	27.7-37.1	236	23.7	19.8-27.6	197	22.4	18.7-26.1	85	12.2	08.9-15.5	62	9.3	06.4-12.2
Male	358	71,708	111	29.4	22.3-36.5	103	25.0	18.9-31.1	77	23.0	16.9-29.1	42	13.3	08.4-18.2	25	9.3	05.2-13.4
Female	541	81,814	208	35.0	28.7-41.3	133	22.6	17.7-27.5	120	21.9	17.4-26.4	43	11.2	06.9-15.5	37	9.3	05.4-13.2
Non-Native	856	142,508	308	31.7	27.0-36.4	221	24.1	20.0-28.2	188	22.7	18.8-26.6	82	12.6	09.1-16.1	57	8.9	06.2-11.6
25-34	208	46,862	9	4.0	01.3-06.7	32	15.5	09.8-21.2	81	40.4	32.8-48.0	46	21.4	15.5-27.3	40	18.8	12.7-24.9
35-44	345	46,572	98	28.5	23.2-33.8	124	38.3	32.2-44.4	86	22.0	17.3-26.7	24	7.0	04.1-09.9	13	4.2	01.7-06.7
45-54	247	25,409	162	64.5	57.8-71.2	53	19.9	14.6-25.2	23	10.8	06.3-15.3	6	3.3	00.6-06.0	3	1.4	00.0-03.2
55-64	57	4,742	30	51.5	37.2-65.8	18	30.7	18.0-43.4	3	4.0	00.0-08.7	4	10.5	00.0-21.1	2	3.3	00.0-07.8
H.S. or G.E.D.	178	38,282	63	31.4	21.0-41.8	54	21.4	14.3-28.5	32	23.3	13.9-32.7	22	17.8	08.2-27.4	7	6.1	01.4-10.8
Some Post-H.S.	280	60,336	111	36.2	28.2-44.2	66	24.1	16.8-31.4	50	17.6	12.3-22.9	31	11.4	07.3-15.5	22	10.7	05.6-15.8
College Graduate	408	41,926	131	24.8	20.5-29.1	107	24.9	20.2-29.6	107	29.3	24.2-34.4	31	10.4	06.5-14.3	32	10.7	06.6-14.8
\$15,000- 24,999	50	12,315	16	26.0	08.8-43.2	11	16.8	03.9-29.7	15	38.2	20.0-56.4	5	11.7	00.0-23.5	3	7.3	00.0-16.5
\$25,000- 34,999	71	13,405	30	24.3	13.3-35.3	14	17.5	06.9-28.1	14	27.1	12.0-42.2	6	14.7	00.0-31.9	7	16.4	04.2-28.6
\$35,000- 49,999	105	19,676	35	31.8	18.1-45.5	26	20.6	11.6-29.6	24	21.0	11.8-30.2	11	12.5	04.9-20.1	9	14.1	03.3-24.9
\$50,000- 74,999	213	35,802	77	34.2	24.6-43.8	53	22.5	15.4-29.6	38	19.0	12.3-25.7	27	14.8	09.1-20.5	18	9.5	04.6-14.4
\$75,000+	383	48,305	134	32.7	26.2-39.2	111	26.7	21.6-31.8	87	23.7	18.6-28.8	33	11.9	07.4-16.4	18	5.0	02.5-07.5

^{&#}x27;% = Percentage, C.I.(95%) = Confidence Interval (at 95 percent probability level). Percentages are weighted to population characteristics.

Rows with less than 50 observations have been suppressed.

TABLE 2: WHAT IS THE BIRTH MONTH AND YEAR OF THE "XTH" CHILD? (RCSBIRTH)

Denominator excludes: Respondents who are female or respondents with do not know/refused or with missing responses

Note: The tables for RCSBIRTH were produced using _CHILDWT as the weighting variable

Denominator excludes respondents not reporting children

Data was weighted using _LANDWT.

DEMOGRAPHIC	RESPONDE	NT NUMBER		1993-1	997		1998-2	002		2003-2	007		2008-2	009		2010-2	011
GROUPS	TOTAL	WEIGHTED	N	%	C.I. (95%)	N	%	C.I. (95%)	N	%	C.I. (95%)	N	%	C.I. (95%)	N	%	C.I. (95%)
Male	358	71,708	111	29.4	22.3-36.5	103	25.0	18.9-31.1	77	23.0	16.9-29.1	42	13.3	08.4-18.2	25	9.3	05.0-13.6
Non-Native	345	67,527	108	29.1	22.0-36.2	98	25.3	19.0-31.6	75	23.9	17.6-30.2	40	13.4	08.3-18.5	24	8.4	04.9-11.9
25-34	71	20,269	2	2.2	00.0-05.3	9	16.1	05.7-26.5	26	38.0	25.5-50.5	19	23.7	13.5-33.9	15	19.9	10.1-29.7
35-44	131	21,826	24	19.5	11.7-27.3	51	40.7	30.9-50.5	36	23.0	15.6-30.4	15	11.4	05.5-17.3	5	5.5	00.6-10.4
45-54	102	13,000	57	56.7	46.1-67.3	28	25.2	16.4-34.0	10	10.8	03.9-17.7	4	4.5	00.0-09.0	3	2.7	00.0-06.0
H.S. or G.E.D.	82	19,817	23	27.1	12.4-41.8	25	18.2	09.8-26.6	16	29.1	14.2-44.0	14	19.4	06.5-32.3	4	6.3	00.0-12.6
Some Post-H.S.	110	27,885	40	32.8	20.8-44.8	30	29.1	17.1-41.1	14	11.6	05.1-18.1	14	13.2	06.3-20.1	12	13.3	04.7-21.9
College Graduate	149	17,621	40	20.4	13.9-26.9	43	25.4	18.0-32.8	43	33.1	24.5-41.7	14	11.5	04.8-18.2	9	9.6	02.5-16.7
\$50,000- 74,999	97	21,035	32	32.9	19.0-46.8	22	20.9	10.9-30.9	18	20.4	10.4-30.4	19	18.6	10.0-27.2	6	7.2	01.1-13.3
\$75,000+	158	22,716	46	29.8	19.6-40.0	53	28.0	20.2-35.8	37	24.7	16.7-32.7	15	12.3	05.8-18.8	7	5.2	01.1-09.3

^{&#}x27;% = Percentage, C.I.(95%) = Confidence Interval (at 95 percent probability level). Percentages are weighted to population characteristics.

Rows with less than 50 observations have been suppressed.

TABLE 3: WHAT IS THE BIRTH MONTH AND YEAR OF THE "XTH" CHILD? (RCSBIRTH)

Denominator excludes: Respondents who are male or respondents with do not know/refused or with missing responses

Note: The tables for RCSBIRTH were produced using _CHILDWT as the weighting variable

Denominator excludes respondents not reporting children

Data was weighted using _LANDWT.

DEMOGRAPHIC	RESPONDE	NT NUMBER		1993-1	997		1998-2	002		2003-2	007		2008-2	009		2010-2	011
GROUPS	TOTAL	WEIGHTED	N	%	C.I. (95%)	N	%	C.I. (95%)	N	%	C.I. (95%)	N	%	C.I. (95%)	N	%	C.I. (95%)
Female	541	81,814	208	35.0	28.7-41.3	133	22.6	17.7-27.5	120	21.9	17.4-26.4	43	11.2	06.9-15.5	37	9.3	05.4-13.2
Non-Native	511	74,980	200	34.1	27.8-40.4	123	23.1	17.8-28.4	113	21.6	17.1-26.1	42	11.9	07.2-16.6	33	9.3	05.2-13.4
25-34	137	26,593	7	5.4	01.3-09.5	23	15.0	08.9-21.1	55	42.2	32.6-51.8	27	19.6	12.5-26.7	25	17.9	10.5-25.3
35-44	214	24,746	74	36.5	29.2-43.8	73	36.2	28.9-43.5	50	21.1	15.2-27.0	9	3.2	01.0-05.4	8	3.1	00.9-05.3
45-54	145	12,409	105	72.7	64.9-80.5	25	14.4	08.7-20.1	13	10.8	05.1-16.5	2	2.1	00.0-05.2			
H.S. or G.E.D.	96	18,464	40	36.1	21.2-51.0	29	24.8	13.6-36.0	16	17.1	07.3-26.9	8	16.1	02.0-30.2	3	5.9	00.0-13.0
Some Post-H.S.	170	32,451	71	39.1	28.5-49.7	36	19.9	11.7-28.1	36	22.8	15.0-30.6	17	9.8	04.9-14.7	10	8.5	02.8-14.2
College Graduate	259	24,305	91	28.0	22.3-33.7	64	24.5	18.6-30.4	64	26.5	20.6-32.4	17	9.5	04.8-14.2	23	11.4	06.5-16.3
\$35,000- 49,999	65	10,425	20	32.0	14.4-49.6	16	20.7	09.7-31.7	17	25.8	13.1-38.5	9	17.7	06.3-29.1	3	3.7	00.0-08.2
\$50,000- 74,999	116	14,767	45	35.9	23.9-47.9	31	24.8	15.6-34.0	20	17.0	09.4-24.6	8	9.4	02.5-16.3	12	12.8	05.0-20.6
\$75,000+	225	25,589	88	35.3	26.9-43.7	58	25.6	18.7-32.5	50	22.8	16.3-29.3	18	11.5	05.4-17.6	11	4.8	01.9-07.7

^{&#}x27;% = Percentage, C.I.(95%) = Confidence Interval (at 95 percent probability level). Percentages are weighted to population characteristics.

Rows with less than 50 observations have been suppressed.

TABLE 4: IS THE CHILD A BOY OR A GIRL? (RCSGENDR)

Denominator excludes: Respondents with do not know/refused/missing responses

Note: The tables for RCSGENDR were produced using _CHILDWT as the weighting variable

Denominator excludes respondents not reporting children

DEMOGRAPHIC	PESPONDE	NT NUMBER		Boy Male			Gir Fema	
GROUPS	TOTAL	WEIGHTED	N	%	C.I. (95%)	N	%	C.I. (95%)
TOTAL	956	165,514	492	50.4	45.7-55.1	464	49.6	44.9-54.3
Male	391	79,067	205	48.1	41.0-55.2	186	51.9	44.8-59.0
Female	565	86,447	287	52.4	46.1-58.7	278	47.6	41.3-53.9
Non-Native	909	151,891	466	51.0	46.3-55.7	443	49.0	44.3-53.7
25-34	211	47,574	107	50.8	43.2-58.4	104	49.2	41.6-56.8
35-44	361	51,095	184	49.2	42.9-55.5	177	50.8	44.5-57.1
45-54	271	28,389	142	54.4	47.7-61.1	129	45.6	38.9-52.3
55-64	64	5,243	34	51.4	37.9-64.9	30	48.6	35.1-62.1
H.S. or G.E.D.	194	42,639	103	49.6	39.2-60.0	91	50.4	40.0-60.8
Some Post-H.S.	294	65,116	144	47.3	39.5-55.1	150	52.7	44.9-60.5
College Graduate	430	43,866	223	53.2	47.9-58.5	207	46.8	41.5-52.1
\$15,000- 24,999	51	12,449	27	49.6	31.0-68.2	24	50.4	31.8-69.0
\$25,000- 34,999	74	13,211	41	48.9	33.2-64.6	33	51.1	35.4-66.8
\$35,000- 49,999	114	20,851	57	51.6	38.9-64.3	57	48.4	35.7-61.1
\$50,000- 74,999	229	38,530	114	47.5	38.7-56.3	115	52.5	43.7-61.3
\$75,000+	398	51,663	200	46.6	40.1-53.1	198	53.4	46.9-59.9

NOTE:

^{&#}x27;% = Percentage, C.I.(95%) = Confidence Interval (at 95 percent probability level). Percentages are weighted to population characteristics.

Rows with less than 50 observations have been suppressed.

TABLE 5: IS THE CHILD A BOY OR A GIRL? (RCSGENDR)

Denominator excludes: Respondents who are female or respondents with do not know/refused or with missing responses

Note: The tables for RCSGENDR were produced using _CHILDWT as the weighting variable

Denominator excludes respondents not reporting children

Data was weighted using _LANDWT.

DEMOGRAPHIC	RESPONDE	NT NUMBER		Boy Male		Girl Female				
GROUPS	TOTAL	WEIGHTED	N	%	C.I. (95%)	N	%	C.I. (95%)		
Male	391	79,067	205	48.1	41.0-55.2	186	51.9	44.8-59.0		
Non-Native	375	72,480	197	49.7	42.6-56.8	178	50.3	43.2-57.4		
25-34	73	20,774	36	52.2	39.7-64.7	37	47.8	35.3-60.3		
35-44	139	25,426	70	45.4	35.2-55.6	69	54.6	44.4-64.8		
45-54	116	15,028	66	57.7	47.7-67.7	50	42.3	32.3-52.3		
H.S. or G.E.D.	97	23,984	47	46.3	32.0-60.6	50	53.7	39.4-68.0		
Some Post-H.S.	117	30,216	57	44.4	32.4-56.4	60	55.6	43.6-67.6		
College Graduate	159	18,711	88	54.4	45.6-63.2	71	45.6	36.8-54.4		
\$50,000- 74,999	107	22,998	55	44.5	32.2-56.8	52	55.5	43.2-67.8		
\$75,000+	169	24,341	90	52.1	42.7-61.5	79	47.9	38.5-57.3		

July 26, 2012

NOTE:

^{&#}x27;% = Percentage, C.I.(95%) = Confidence Interval (at 95 percent probability level). Percentages are weighted to population characteristics.

Rows with less than 50 observations have been suppressed.

TABLE 6: IS THE CHILD A BOY OR A GIRL? (RCSGENDR)

Denominator excludes: Respondents who are male or respondents with do not know/refused or with missing responses

Note: The tables for RCSGENDR were produced using _CHILDWT as the weighting variable

Denominator excludes respondents not reporting children

Data was weighted using _LANDWT.

DEMOGRAPHIC	RESPONDE	NT NUMBER		Boy Mal			Gir Fema	
GROUPS	TOTAL	WEIGHTED	N	%	C.I. (95%)	N	%	C.I. (95%)
Female	565	86,447	287	52.4	46.1-58.7	278	47.6	41.3-53.9
Non-Native	534	79,411	269	52.2	45.7-58.7	265	47.8	41.3-54.3
25-34	138	26,801	71	49.7	40.3-59.1	67	50.3	40.9-59.7
35-44	222	25,670	114	53.0	45.7-60.3	108	47.0	39.7-54.3
45-54	155	13,360	76	50.7	42.1-59.3	79	49.3	40.7-57.9
H.S. or G.E.D.	97	18,655	56	53.8	38.7-68.9	41	46.2	31.1-61.3
Some Post-H.S.	177	34,900	87	49.8	39.2-60.4	90	50.2	39.6-60.8
College Graduate	271	25,155	135	52.3	45.6-59.0	136	47.7	41.0-54.4
\$35,000- 49,999	69	10,824	36	56.8	41.9-71.7	33	43.2	28.3-58.1
\$50,000- 74,999	122	15,531	59	52.0	40.8-63.2	63	48.0	36.8-59.2
\$75,000+	229	27,322	110	41.7	33.1-50.3	119	58.3	49.7-66.9

NOTE:

^{&#}x27;% = Percentage, C.I.(95%) = Confidence Interval (at 95 percent probability level). Percentages are weighted to population characteristics.

Rows with less than 50 observations have been suppressed.

TABLE 7. IS THE CHILD HISPANIC OR LATINO? (RCHISLAT)

Denominator excludes: Respondents with do not know/refused/missing responses

Note: The tables for RCHISLAT were produced using _CHILDWT as the weighting variable

Denominator excludes respondents not reporting children

Data was weighted using LANDWT.

DEMOGRAPHIC	RESPONDE	as weighted NT NUMBER	usin	y Yes	indwi.		No	
GROUPS	TOTAL	WEIGHTED	N	%	C.I. (95%)	N	%	C.I. (95%)
TOTAL	960	166,400	26	3.7	01.9-05.5	934	96.3	94.5-98.1
Male	393	79,550	11	3.1	01.1-05.1	382	96.9	94.9-98.9
Female	567	86,850	15	4.4	01.5-07.3	552	95.6	92.7-98.5
Non-Native	914	152,891	26	4.1	02.1-06.1	888	95.9	93.9-97.9
25-34	212	47,782	7	3.5	00.8-06.2	205	96.5	93.8-99.2
35-44	363	51,482	9	4.1	01.2-07.0	354	95.9	93.0-98.8
45-54	271	28,617	5	2.0	00.0-04.0	266	98.0	96.0-0100
55-64	65	5,305	3	6.5	00.0-13.6	62	93.5	86.4-0100
H.S. or G.E.D.	193	42,567	6	2.8	00.4-05.2	187	97.2	94.8-99.6
Some Post-H.S.	298	66,050	7	2.2	00.4-04.0	291	97.8	96.0-99.6
College Graduate	431	43,890	8	2.3	00.5-04.1	423	97.7	95.9-99.5
\$15,000- 24,999	50	12,377	2	4.6	00.0-11.9	48	95.4	88.1-0100
\$25,000- 34,999	74	13,211	4	7.1	00.0-14.9	70	92.9	85.1-0100
\$35,000- 49,999	113	20,720	6	5.8	00.9-10.7	107	94.2	89.3-99.1
\$50,000- 74,999	230	38,726	5	2.5	00.1-04.9	225	97.5	95.1-99.9
\$75,000+	401	52,258	5	1.6	00.0-03.2	396	98.4	96.8-0100

July 26, 2012

NOTE:

^{&#}x27;% = Percentage, C.I.(95%) = Confidence Interval (at 95 percent probability level). Percentages are weighted to population characteristics.

Rows with less than 50 observations have been suppressed.

TABLE 8: IS THE CHILD HISPANIC OR LATINO? (RCHISLAT)

Denominator excludes: Respondents who are female or respondents with do not know/refused or with missing responses

Note: The tables for RCHISLAT were produced using _CHILDWT as the weighting variable

Denominator excludes respondents not reporting children

Data was weighted using _LANDWT.

				J				
DEMOGRAPHIC	RESPONDE	NT NUMBER		Yes	3		No	,
GROUPS	TOTAL	WEIGHTED	N	%	C.I. (95%)	N	%	C.I. (95%)
Male	393	79,550	11	3.1	01.1-05.1	382	96.9	94.9-98.9
Non-Native	378	73,077	11	3.3	01.1-05.5	367	96.7	94.5-98.9
25-34	73	20,774				73	100.0	0100-0100
35-44	140	25,608	7	7.6	01.9-13.3	133	92.4	86.7-98.1
45-54	117	15,329	3	2.5	00.0-05.4	114	97.5	94.6-0100
H.S. or G.E.D.	97	23,984	5	4.5	00.4-08.6	92	95.5	91.4-99.6
Some Post-H.S.	119	30,738	2	1.6	00.0-04.0	117	98.4	96.0-0100
College Graduate	159	18,672	3	2.0	00.0-04.4	156	98.0	95.6-0100
\$50,000- 74,999	108	23,195	2	2.1	00.0-05.0	106	97.9	95.0-0100
\$75.000+	170	24.523	2	1.6	00.0-04.0	168	98.4	96.0-0100

^{&#}x27;% = Percentage, C.I.(95%) = Confidence Interval (at 95 percent probability level). Percentages are weighted to population characteristics.

Rows with less than 50 observations have been suppressed.

TABLE 9: IS THE CHILD HISPANIC OR LATINO? (RCHISLAT)

Denominator excludes: Respondents who are male or respondents with do not know/refused or with missing responses
Note: The tables for RCHISLAT were produced using _CHILDWT as the weighting variable

Denominator excludes respondents not reporting children

Data was weighted using _LANDWT.

	Dava II	ab "CI31100	~ ~~===	J				
DEMOGRAPHIC	RESPONDE	NT NUMBER		Yes	3		No	
GROUPS	TOTAL	WEIGHTED	N	%	C.I. (95%)	N	%	C.I. (95%)
Female	567	86,850	15	4.4	01.5-07.3	552	95.6	92.7-98.5
Non-Native	536	79,813	15	4.7	01.6-07.8	521	95.3	92.2-98.4
25-34	139	27,009	7	6.2	01.3-11.1	132	93.8	88.9-98.7
35-44	223	25,874	2	0.7	00.0-01.7	221	99.3	98.3-0100
45-54	154	13,288	2	1.3	00.0-03.3	152	98.7	96.7-0100
H.S. or G.E.D.	96	18,583	1	0.7	00.0-02.1	95	99.3	97.9-0100
Some Post-H.S.	179	35,312	5	2.7	00.2-05.2	174	97.3	94.8-99.8
College Graduate	272	25,217	5	2.4	00.0-04.9	267	97.6	95.1-0100
\$35,000- 49,999	69	10,824	1	1.9	00.0-05.6	68	98.1	94.4-0100
\$50,000- 74,999	122	15,531	3	3.1	00.0-07.2	119	96.9	92.8-0100
\$75,000+	231	27 734	3	16	00 0-03 6	228	98 4	96 4-0100

^{&#}x27;% = Percentage, C.I.(95%) = Confidence Interval (at 95 percent probability level). Percentages are weighted to population characteristics.

Rows with less than 50 observations have been suppressed.

TABLE 10: WHICH ONE OR MORE OF THE FOLLOWING WOULD YOU SAY IS THE RACE OF THE CHILD? (RCSRACE)

Denominator excludes: Respondents with do not know/refused/missing responses

Note: The tables for RCSRACE were produced using _CHILDWT as the weighting variable

Denominator excludes respondents not reporting children

Data was weighted using _LANDWT.

Am. Indian or Alaskan **DEMOGRAPHIC** RESPONDENT NUMBER White **Black or African American** Asian Native Hawaiian or P.I. Native Other **GROUPS TOTAL** WEIGHTED Ν C.I. (95%) C.I. (95%) Ν % C.I. (95%) C.I. (95%) C.I. (95%) Ν C.I. (95%) TOTAL 950 164,179 857 85.7 82.0-89.4 9 1.1 00.3-01.9 12 1.5 00.3-02.7 2 0.0 00.0-00.0 53 9.8 06.3-13.3 17 1.9 00.7-03.1 77,956 86.4 80.3-92.5 1.2 00.0-02.8 00.4-02.8 04.1-15.5 00.0-02.8 Male 355 1.6 19 9.8 2 1.0 00.0-01.9 Female 563 86.223 502 85.1 80.4-89.8 5 0.9 5 1.5 00.0-03.5 2 0.1 00.0-00.3 34 9.8 05.5-14.1 15 2.6 01.0-04.2 Non-Native 905 150,830 851 91.9 89.0-94.8 9 1.2 00.2-02.2 12 1.7 00.5-02.9 0.1 00.1-00.1 16 3.3 00.9-05.7 15 1.9 00.5-03.3 25-34 211 47,570 186 85.3 79.4-91.2 2 1.5 00.0-03.9 4 1.4 00.0-02.8 14 8.8 04.1-13.5 2.9 00.0-06.2 5 35-44 357 50,031 327 86.4 80.3-92.5 5 00.0-02.8 5 1.5 00.1-02.9 16 8.6 02.9-14.3 2.1 00.0-04.3 45-54 269 28.201 249 91.4 87.3-95.5 2 1.1 00.0-02.7 0.9 00.0-02.3 0.3 00.0-00.7 10 5.1 01.6-08.6 1.2 00.0-02.4 55-64 64 5.165 56 84.8 74.0-95.6 5 11.2 01.0-21.4 3 4.0 00.0-08.7 H.S. or G.E.D. 78.2-93.4 0.4 00.0-01.2 0.6 00.0-01.8 04.4-19.2 00.0-03.2 190 41,881 172 85.8 13 11.8 1.4 Some Post-H.S. 296 65.567 271 86.9 80.2-93.6 2 0.7 00.0-01.7 18 10.0 03.7-16.3 5 2.4 00.0-04.9 0.2 00.0-00.4 00.0-05.4 College Graduate 428 43,457 389 88.5 84.6-92.4 6 2.7 10 3.2 01.0-05.4 15 4.0 01.8-06.2 6 1.3 00.1-02.5 \$15,000-24,999 12,236 43.8-79.4 00.0-18.8 06.0-33.4 00.0-21.9 50 35 61.6 8 19.7 9.6 \$25,000-34,999 72 12,606 63 87.1 78.3-95.9 2 00.0-05.7 5 8.2 00.6-15.8 2 2.3 00.0-05.6 \$35.000-49.999 113 20.590 106 91.0 81.8-0100 1.3 00.0-03.8 5 7.1 00.0-16.1 0.6 00.0-01.8

0.7 00.0-01.9

00.0-00.6

2.3 00.7-03.9

\$50.000-74.999

\$75,000+

229

397

38.503

51,610

217

94.6

93.6

91.1-98.1

90.9-96.3

NOTE:

Cells that contain a confidence interval with a half-width of greater than 10 have been suppressed.'

5

12

0.2 00.0-00.4

2.6

00.1-05.1

01.1-05.1

5

2.1

0.7

00.0-04.5

00.0-01.5

^{&#}x27;% = Percentage, C.I.(95%) = Confidence Interval (at 95 percent probability level). Percentages are weighted to population characteristics.

Rows with less than 50 observations have been suppressed.

BEHAVIORAL RISK FACTOR SURVEILLANCE SYSTEM NORTH DAKOTA MODULE QUESTIONS DATA REPORT, 2011

MODULE 32: RANDOM CHILD SELECTION

TABLE 11: WHICH ONE OR MORE OF THE FOLLOWING WOULD YOU SAY IS THE RACE OF THE CHILD? (RCSRACE)

Denominator excludes: Respondents who are female or respondents with do not know/refused or with missing responses Note: The tables for RCSRACE were produced using _CHILDWT as the weighting variable

Denominator excludes respondents not reporting children

Data was weighted using _LANDWT.

DEMOGRAPHIC	RESPONDE	NT NUMBER		Whit	е	Black o	r Africa	n American		Asia	n	Am. Ind	ian or Al	askan Native		Othe	r
GROUPS	TOTAL	WEIGHTED	N	%	C.I. (95%)	N	%	C.I. (95%)	N	%	C.I. (95%)	N	%	C.I. (95%)	N	%	C.I. (95%)
Male	387	77,956	355	86.4	80.3-92.5	4	1.2	00.0-02.8	7	1.6	00.4-02.8	19	9.8	04.1-15.5	2	1.0	00.0-02.8
Non-Native	373	71,643	352	91.6	86.7-96.5	4	1.4	00.0-03.0	7	1.7	00.3-03.1	8	4.3	00.0-08.6	2	1.1	00.0-03.1
25-34	73	20,774	66	87.3	77.7-96.9	1	2.5	00.0-07.4	2	1.8	00.0-04.3	3	4.8	00.0-10.3	1	3.5	00.0-10.4
35-44	134	24,157	124	84.8	73.8-95.8	2	1.3	00.0-03.1	3	2.4	00.0-05.1	5	11.5	00.5-22.5			
45-54	117	15,186	108	90.2	83.5-96.9	1	0.9	00.0-02.7	2	1.7	00.0-04.2	6	7.2	01.3-13.1			
H.S. or G.E.D.	94	23,352	87	90.3	82.1-98.5	1	0.6	00.0-01.8	1	1.1	00.0-03.3	5	8.0	00.2-15.8			
Some Post-H.S.	118	30,413	112	83.6	70.9-96.3							5	14.0	01.7-26.3	1	2.4	00.0-07.1
College Graduate	158	18,513	143	87.4	80.1-94.7	3	4.4	00.0-10.3	6	5.1	01.0-09.2	5	2.8	00.1-05.5	1	0.3	00.0-00.9
\$50,000- 74,999	107	22,972	104	96.9	92.8-0100							2	2.9	00.0-07.0	1	0.2	00.0-00.6
\$75,000+	167	24,034	158	95.1	91.6-98.6				5	3.0	00.3-05.7	4	1.9	00.0-04.3			

^{&#}x27;% = Percentage, C.I.(95%) = Confidence Interval (at 95 percent probability level). Percentages are weighted to population characteristics. Rows with less than 50 observations have been suppressed.

BEHAVIORAL RISK FACTOR SURVEILLANCE SYSTEM NORTH DAKOTA MODULE QUESTIONS DATA REPORT, 2011

MODULE 32: RANDOM CHILD SELECTION

TABLE 12: WHICH ONE OR MORE OF THE FOLLOWING WOULD YOU SAY IS THE RACE OF THE CHILD? (RCSRACE)

Denominator excludes: Respondents who are male or respondents with do not know/refused or with missing responses Note: The tables for RCSRACE were produced using _CHILDWT as the weighting variable

Denominator excludes respondents not reporting children

Data was weighted using _LANDWT.

DEMOGRAPHIC	RESPONDE	NT NUMBER		Whi	te	Black or	r Africa	n American		Asia	ın	Native	Hawa	iian or P.I.	Am. I	ndian o Nativ	r Alaskan ⁄e		Othe	er er
GROUPS	TOTAL	WEIGHTED	N	%	C.I. (95%)	N	%	C.I. (95%)	N	%	C.I. (95%)	N	%	C.I. (95%)	N	%	C.I. (95%)	N	%	C.I. (95%)
Female	563	86,223	502	85.1	80.4-89.8	5	0.9	00.0-01.9	5	1.5	00.0-03.5	2	0.1	00.0-00.3	34	9.8	05.7-13.9	15	2.6	01.0-04.2
Non-Native	532	79,187	499	92.2	88.7-95.7	5	1.0	00.0-02.0	5	1.6	00.0-03.8	2	0.1	00.0-00.3	8	2.5	00.3-04.7	13	2.6	01.0-04.2
25-34	138	26,796	120	83.8	76.4-91.2	1	0.8	00.0-02.4	2	1.1	00.0-02.7				11	11.9	04.8-19.0	4	2.4	00.0-04.8
35-44	223	25,874	203	87.9	82.4-93.4	3	1.6	00.0-03.8	2	0.6	00.0-01.4				11	5.9	02.4-09.4	4	4.0	00.0-08.1
45-54	152	13,015	141	92.7	88.0-97.4	1	1.4	00.0-04.1				2	0.6	00.0-01.6	4	2.7	00.0-05.4	4	2.5	00.0-05.0
H.S. or G.E.D.	96	18,528	85	80.1	67.0-93.2										8	16.7	03.8-29.6	3	3.2	00.0-07.3
Some Post-H.S.	178	35,154	159	89.8	84.9-94.7	2	1.3	00.0-03.1							13	6.5	02.6-10.4	4	2.4	00.0-04.9
College Graduate	270	24,944	246	89.4	85.1-93.7	3	1.4	00.0-03.2	4	1.8	00.0-03.6	2	0.3	00.0-00.7	10	4.9	01.6-08.2	5	2.1	00.1-04.1
\$35,000- 49,999	69	10,824	66	96.8	93.1-0100										2	2.1	00.0-05.0	1	1.1	00.0-03.3
\$50,000-74,999	122	15,531	113	91.4	84.9-97.9	2	1.8	00.0-04.5							3	2.1	00.0-04.5	4	4.8	00.0-10.3
\$75,000+	230	27,576	212	92.4	88.5-96.3	1	0.3	00.0-00.9	4	1.7	00.0-03.5	2	0.3	00.0-00.7	8	4.1	01.0-07.2	3	1.2	00.0-02.8

NOTE:

Ann Indian an Alaskan

^{&#}x27;% = Percentage, C.I.(95%) = Confidence Interval (at 95 percent probability level). Percentages are weighted to population characteristics. Rows with less than 50 observations have been suppressed.

TABLE 13: WHICH ONE OF THESE GROUPS WOULD YOU SAY BEST REPRESENTS THE CHILD'S RACE? (RCSBRACE)

Denominator excludes: Respondents with do not know/refused/missing responses

Note: The tables for RCSBRACE were produced using _CHILDWT as the weighting variable

Denominator excludes respondents not reporting multiple race groups

There is not enough data (less than 50 total observations) to print this table.

NOTE:

'% = Percentage, C.I.(95%) = Confidence Interval (at 95 percent probability level). Percentages are weighted to population characteristics.

Rows with less than 50 observations have been suppressed.

TABLE 14: WHICH ONE OF THESE GROUPS WOULD YOU SAY BEST REPRESENTS THE CHILD'S RACE? (RCSBRACE)

Denominator excludes: Respondents who are female or respondents with do not know/refused or with missing responses

Note: The tables for RCSBRACE were produced using _CHILDWT as the weighting variable

Denominator excludes respondents not reporting multiple race groups

There is not enough data (less than 50 total observations) to print this table.

NOTE:

'% = Percentage, C.I.(95%) = Confidence Interval (at 95 percent probability level). Percentages are weighted to population characteristics.

Rows with less than 50 observations have been suppressed.

TABLE 15: WHICH ONE OF THESE GROUPS WOULD YOU SAY BEST REPRESENTS THE CHILD'S RACE? (RCSBRACE)

Denominator excludes: Respondents who are male or respondents with do not know/refused or with missing responses

Note: The tables for RCSBRACE were produced using $_$ CHILDWT as the weighting variable

Denominator excludes respondents not reporting multiple race groups

There is not enough data (less than 50 total observations) to print this table.

NOTE:

'% = Percentage, C.I.(95%) = Confidence Interval (at 95 percent probability level). Percentages are weighted to population characteristics.

Rows with less than 50 observations have been suppressed.

TABLE 16: HOW ARE YOU RELATED TO THE CHILD? (RCSRLTN2)

Denominator excludes: Respondents with do not know/refused/missing responses Note: The tables for RCSRLTN2 were produced using _CHILDWT as the weighting variable

Data was weighted using _LANDWT.

									F	oster p	parent									
DEMOGRAPHIC	RESPONDE	NT NUMBER		Pare	ent	(Grandp	arent		guard	lian		Sibli	ng	С	ther re	elative		Not rel	ated
GROUPS	TOTAL	WEIGHTED	N	%	C.I. (95%)	N	%	C.I. (95%)	N	%	C.I. (95%)	N	%	C.I. (95%)	N	%	C.I. (95%)	N	%	C.I. (95%)
TOTAL	966	166,916	890	78.6	73.1-84.1	31	2.2	01.4-03.0	10	1.1	00.3-01.9	19	12.5	07.4-17.6	9	4.0	01.1-06.9	7	1.6	00.0-03.2
Male	395	79,563	358	76.4	68.2-84.6	11	1.8	00.6-03.0	9	2.2	00.6-03.8	9	13.1	05.5-20.7	5	5.3	00.4-10.2	3	1.2	00.0-03.0
Female	571	87,354	532	80.6	73.3-87.9	20	2.5	01.3-03.7	1	0.2	00.0-00.6	10	11.9	05.2-18.6	4	2.8	00.0-06.3	4	2.1	00.0-04.5
Non-Native	918	153,185	857	81.2	75.7-86.7	27	2.0	01.2-02.8	6	0.6	00.0-01.2	16	11.5	06.4-16.6	5	3.0	00.3-05.7	7	1.8	00.2-03.4
25-34	214	48,171	208	95.0	90.5-99.5				1	0.3	00.0-00.9				3	3.8	00.0-08.3	2	0.9	00.0-02.1
35-44	363	51,185	356	93.2	87.5-98.9				3	1.5	00.0-03.3				3	3.9	00.0-09.0	1	1.4	00.0-03.9
45-54	274	28,935	260	92.8	88.9-96.7	7	3.6	00.9-06.3	6	3.3	00.4-06.2							1	0.2	00.0-00.6
55-64	65	5,305	48	70.6	58.1-83.1	15	26.5	14.2-38.8							1	1.7	00.0-05.0	1	1.3	00.0-03.8
H.S. or G.E.D.	194	42,342	170	73.0	61.2-84.8	9	2.1	00.7-03.5	2	0.8	00.0-02.0	7	17.6	06.4-28.8	4	5.8	00.0-12.7	2	0.7	00.0-01.9
Some Post-H.S.	300	66,288	274	75.5	66.1-84.9	9	1.8	00.4-03.2	3	1.2	00.0-02.6	8	14.1	05.5-22.7	5	6.3	00.4-12.2	1	1.0	00.0-03.0
College Graduate	434	44,393	422	97.4	95.8-99.0	7	1.5	00.3-02.7	4	1.0	00.0-02.0							1	0.1	00.0-00.3
\$15,000- 24,999	52	12,676	41	69.4	50.0-88.8	6	7.2	00.7-13.7	2	2.4	00.0-05.7	1	8.9	00.0-25.2	2	12.0	00.0-27.3			
\$25,000- 34,999	73	12,817	66	91.3	83.9-98.7	4	3.5	00.0-07.0	1	2.3	00.0-06.6				1	2.5	00.0-07.4	1	0.5	00.0-01.5
\$35,000- 49,999	114	20,851	109	86.8	72.7-0100	3	1.9	00.0-04.3				2	11.3	00.0-25.4						
\$50,000- 74,999	230	38,726	216	85.8	76.2-95.4	6	1.8	00.2-03.4	1	0.5	00.0-01.5	3	8.8	00.0-18.0				4	3.0	00.0-06.7
\$75,000+	402	52,419	387	90.3	82.9-97.7	8	1.5	00.5-02.5	4	1.2	00.0-02.4	2	4.4	00.0-10.3	1	2.7	00.0-07.8			

NOTE:

^{&#}x27;% = Percentage, C.I.(95%) = Confidence Interval (at 95 percent probability level). Percentages are weighted to population characteristics.

Rows with less than 50 observations have been suppressed.

TABLE 17: HOW ARE YOU RELATED TO THE CHILD? (RCSRLTN2)

Denominator excludes: Respondents who are female or respondents with do not know/refused or with missing responses

Note: The tables for RCSRLTN2 were produced using _CHILDWT as the weighting variable

Data was weighted using _LANDWT.

DEMOGRAPHIC	RESPONDE	NT NUMBER		Pare	ent		Grandp	arent	F	oster p guard			Sibli	na	c	ther re	lative		Not rel	ated
GROUPS	TOTAL	WEIGHTED	N	%	C.I. (95%)	N	%	C.I. (95%)	N	%	C.I. (95%)	N	%	C.I. (95%)	N	%	C.I. (95%)	N	%	C.I. (95%)
Male	395	79,563	358	76.4	68.2-84.6	11	1.8	00.6-03.0	9	2.2	00.6-03.8	9	13.1	05.5-20.7	5	5.3	00.4-10.2	3	1.2	00.0-03.0
Non-Native	379	72,975	348	79.6	71.2-88.0	11	2.0	00.8-03.2	6	1.3	00.1-02.5	8	12.6	04.8-20.4	3	3.2	00.0-07.1	3	1.3	00.0-03.3
25-34	74	20,935	70	93.4	86.0-0100				1	0.8	00.0-02.4				2	5.0	00.0-12.3	1	0.8	00.0-02.4
35-44	139	25,214	134	87.4	76.4-98.4				2	2.3	00.0-05.6				2	7.5	00.0-17.5	1	2.7	0.80-0.00
45-54	119	15,574	111	91.2	84.9-97.5	2	2.6	00.0-06.1	6	6.2	00.7-11.7									
H.S. or G.E.D.	96	23,589	82	67.4	50.9-83.9	4	1.9	00.0-03.9	2	1.5	00.0-03.7	4	18.9	03.2-34.6	3	10.0	00.0-21.8	1	0.3	00.0-00.9
Some Post-H.S.	120	30,869	111	76.1	62.0-90.2	1	0.6	00.0-01.8	2	2.1	00.0-05.0	3	12.9	00.0-25.8	2	6.1	00.0-14.3	1	2.2	00.0-06.5
College Graduate	161	18,948	153	95.5	92.2-98.8	4	2.2	00.0-04.6	4	2.3	00.0-04.7									
\$50,000- 74,999	108	23,195	100	83.2	69.5-96.9	2	1.3	00.0-03.3	1	0.9	00.0-02.7	2	10.5	00.0-23.8				3	4.0	00.0-09.9
\$75,000+	171	24,685	161	90.9	81.9-99.9	5	1.8	00.0-03.6	4	2.5	00.0-05.2	1	4.7	00.0-13.5						

NOTE:

^{&#}x27;% = Percentage, C.I.(95%) = Confidence Interval (at 95 percent probability level). Percentages are weighted to population characteristics.

Rows with less than 50 observations have been suppressed.

TABLE 18: HOW ARE YOU RELATED TO THE CHILD? (RCSRLTN2)

Denominator excludes: Respondents who are male or respondents with do not know/refused or with missing responses

Note: The tables for RCSRLTN2 were produced using _CHILDWT as the weighting variable

Data was weighted using _LANDWT.

DEMOGRAPHIC	RESPONDE	NT NUMBER		Pare	ent	G	randp	arent	F	oster p quard	oarent Jian		Sibli	na	0	ther re	lative		Not rel	ated
GROUPS	TOTAL	WEIGHTED	N	%	C.I. (95%)	N	%	C.I. (95%)	N	%	C.I. (95%)	N	%	C.I. (95%)	N	%	C.I. (95%)	N	%	C.I. (95%)
Female	571	87,354	532	80.6	73.3-87.9	20	2.5	01.3-03.7	1	0.2	00.0-00.6	10	11.9	05.2-18.6	4	2.8	00.0-06.3	4	2.1	00.0-04.5
Non-Native	539	80,210	509	82.6	75.2-90.0	16	2.0	01.0-03.0				8	10.4	03.7-17.1	2	2.7	00.0-06.6	4	2.2	00.0-04.7
25-34	140	27,236	138	96.2	90.5-0100										1	2.9	00.0-08.4	1	0.9	00.0-02.7
35-44	224	25,971	222	98.9	97.3-0100				1	0.6	00.0-01.8				1	0.5	00.0-01.5			
45-54	155	13,360	149	94.7	90.4-99.0	5	4.8	00.7-08.9										1	0.5	00.0-01.5
H.S. or G.E.D.	98	18,753	88	80.0	64.3-95.7	5	2.3	00.1-04.5				3	16.0	00.1-31.9	1	0.5	00.0-01.5	1	1.3	00.0-03.8
Some Post-H.S.	180	35,419	163	74.9	62.4-87.4	8	2.8	00.6-05.0	1	0.5	00.0-01.5	5	15.2	03.6-26.8	3	6.6	00.0-15.0			
College Graduate	273	25,445	269	98.7	97.3-0100	3	1.1	00.0-02.3										1	0.2	00.0-00.6
\$35,000- 49,999	69	10,824	66	87.5	68.9-0100	2	1.8	00.0-04.3				1	10.7	00.0-29.5						
\$50,000- 74,999	122	15,531	116	89.7	77.9-0100	4	2.4	00.0-04.9				1	6.3	00.0-17.9				1	1.5	00.0-04.4
\$75,000+	231	27,734	226	89.8	78.2-0100	3	1.1	00.0-02.5				1	4.0	00.0-11.6	1	5.1	00.0-14.5			

NOTE:

^{&#}x27;% = Percentage, C.I.(95%) = Confidence Interval (at 95 percent probability level). Percentages are weighted to population characteristics.

Rows with less than 50 observations have been suppressed.

BEHAVIORAL RISK FACTOR SURVEILLANCE SYSTEM NORTH DAKOTA MODULE QUESTIONS DATA REPORT, 2011

MODULE 33: CHILDHOOD ASTHMA PREVALENCE CONTENTS

* NOTE	*****************************	* * * * *
WHE	N THE DENOMINATOR IS NOT BASED ON THE TOTAL POPULATION SURVEYED, THE TITLE CONTAINS A DESCRIPTION OF WHAT POPULA	ΓΙΟN
IS E	XCLUDED FROM THE DENOMINATOR. IF THE TITLE DOES NOT DESCRIBE THE DENOMINATOR, THE TABLE IS BASED ON ALL RESPONDE	NTS
	TO THE STATES SURVEY.	
* * * *	*****************************	****
TABLE		PAGE
	HAS A DOCTOR, NURSE OR OTHER HEALTH PROFESSIONAL EVER SAID THAT THE CHILD HAS ASTHMA? (CASTHDX2)	
1	MALES AND FEMALES	86
2	MALES ONLY	87
3	FEMALES ONLY	88
	DOES THE CHILD STILL HAVE ASTHMA? (CASTHNO2)	
4	MALES AND FEMALES	89
5	MALES ONLY	90
6	FEMALES ONLY	91

86 of 93 July 26, 2012

TABLE 1: HAS A DOCTOR, NURSE OR OTHER HEALTH PROFESSIONAL EVER SAID THAT THE CHILD HAS ASTHMA? (CASTHDX2)

Denominator excludes: Respondents with do not know/refused/missing responses

Denominator excludes: Respondents without children

Note: The tables for CASTHDX2 were produced using _CHILDWT as the weighting variable

Data was weighted using LANDWT.

DEMOGRAPHIC	RESPONDE	NT NUMBER		Yes	indwi.		No	
GROUPS	TOTAL	WEIGHTED	N	%	C.I. (95%)	N	%	C.I. (95%)
TOTAL	963	164,052	88	8.0	05.6-10.4	875	92.0	89.6-94.4
Male	392	76,644	29	7.6	03.7-11.5	363	92.4	88.5-96.3
Female	571	87,407	59	8.3	05.8-10.8	512	91.7	89.2-94.2
Non-Native	915	150,321	84	8.3	05.8-10.8	831	91.7	89.2-94.2
25-34	212	47,890	8	6.1	01.6-10.6	204	93.9	89.4-98.4
35-44	364	51,512	42	10.6	07.3-13.9	322	89.4	86.1-92.7
45-54	272	28,560	31	10.6	06.9-14.3	241	89.4	85.7-93.1
55-64	67	5,565	6	8.7	01.1-16.3	61	91.3	83.7-98.9
H.S. or G.E.D.	192	41,049	16	5.4	02.3-08.5	176	94.6	91.5-97.7
Some Post-H.S.	301	64,998	29	9.4	04.7-14.1	272	90.6	85.9-95.3
College Graduate	432	44,112	38	7.7	05.0-10.4	394	92.3	89.6-95.0
\$15,000- 24,999	53	12,765	4	12.6	00.0-25.5	49	87.4	74.5-0100
\$25,000- 34,999	72	12,626	5	4.6	00.0-09.3	67	95.4	90.7-0100
\$35,000- 49,999	115	21,054	11	8.6	03.1-14.1	104	91.4	85.9-96.9
\$50,000- 74,999	229	38,778	25	8.9	04.6-13.2	204	91.1	86.8-95.4
\$75,000+	400	52,034	38	9.6	04.9-14.3	362	90.4	85.7-95.1

NOTE:

^{&#}x27;% = Percentage, C.I.(95%) = Confidence Interval (at 95 percent probability level). Percentages are weighted to population characteristics.

Rows with less than 50 observations have been suppressed.

TABLE 2: HAS A DOCTOR, NURSE OR OTHER HEALTH PROFESSIONAL EVER SAID THAT THE CHILD HAS ASTHMA? (CASTHDX2)

Denominator excludes: Respondents who are female or respondents with do not know/refused or with missing responses

Denominator excludes: Respondents without children

Note: The tables for CASTHDX2 were produced using _CHILDWT as the weighting variable

Data was weighted using _LANDWT.

DEMOGRAPHIC	RESPONDE	NT NUMBER		Yes	3		No)
GROUPS	TOTAL	WEIGHTED	N	%	C.I. (95%)	N	%	C.I. (95%)
Male	392	76,644	29	7.6	03.5-11.7	363	92.4	88.3-96.5
Non-Native	376	70,057	28	8.1	03.8-12.4	348	91.9	87.6-96.2
25-34	73	20,774	2	6.1	00.0-14.1	71	93.9	85.9-0100
35-44	140	25,435	11	6.8	02.5-11.1	129	93.2	88.9-97.5
45-54	118	15,391	12	8.8	03.9-13.7	106	91.2	86.3-96.1
H.S. or G.E.D.	96	22,584	7	4.6	00.9-08.3	89	95.4	91.7-99.1
Some Post-H.S.	118	29,117	12	11.8	02.6-21.0	106	88.2	79.0-97.4
College Graduate	160	18,787	8	3.5	01.0-06.0	152	96.5	94.0-99.0
\$50,000- 74,999	108	23,336	10	7.9	02.0-13.8	98	92.1	86.2-98.0
\$75.000+	170	24.420	9	7.9	00.0-16.9	161	92.1	83.1-0100

NOTE:

July 26, 2012

^{&#}x27;% = Percentage, C.I.(95%) = Confidence Interval (at 95 percent probability level). Percentages are weighted to population characteristics. Rows with less than 50 observations have been suppressed.

BEHAVIORAL RISK FACTOR SURVEILLANCE SYSTEM NORTH DAKOTA MODULE QUESTIONS DATA REPORT. 2011

MODULE 33: CHILDHOOD ASTHMA PREVALENCE

TABLE 3: HAS A DOCTOR, NURSE OR OTHER HEALTH PROFESSIONAL EVER SAID THAT THE CHILD HAS ASTHMA? (CASTHDX2)

Denominator excludes: Respondents who are male or respondents with do not know/refused or with missing responses Denominator excludes: Respondents without children

Note: The tables for CASTHDX2 were produced using _CHILDWT as the weighting variable

Data was weighted using _LANDWT.

	Dava II	Data was werdined asingmmswi.							
DEMOGRAPHIC	RESPONDENT NUMBER			Yes			No		
GROUPS	TOTAL	WEIGHTED	N	%	C.I. (95%)	N	%	C.I. (95%)	
Female	571	87,407	59	8.3	05.8-10.8	512	91.7	89.2-94.2	
Non-Native	539	80,264	56	8.6	05.9-11.3	483	91.4	88.7-94.1	
25-34	139	27,116	6	6.1	01.0-11.2	133	93.9	88.8-99.0	
35-44	224	26,077	31	14.3	09.2-19.4	193	85.7	80.6-90.8	
45-54	154	13,169	19	12.7	07.0-18.4	135	87.3	81.6-93.0	
H.S. or G.E.D.	96	18,464	9	6.3	01.2-11.4	87	93.7	88.6-98.8	
Some Post-H.S.	183	35,881	17	7.4	03.7-11.1	166	92.6	88.9-96.3	
College Graduate	272	25,325	30	10.8	06.7-14.9	242	89.2	85.1-93.3	
\$35,000- 49,999	70	11,026	6	10.1	01.3-18.9	64	89.9	81.1-98.7	
\$50,000- 74,999	121	15,442	15	10.3	04.6-16.0	106	89.7	84.0-95.4	
\$75,000+	230	27,614	29	11.1	06.6-15.6	201	88.9	84.4-93.4	

^{&#}x27;% = Percentage, C.I.(95%) = Confidence Interval (at 95 percent probability level). Percentages are weighted to population characteristics. Rows with less than 50 observations have been suppressed.

TABLE 4: DOES THE CHILD STILL HAVE ASTHMA? (CASTHNO2)

Denominator excludes: Respondents with do not know/refused/missing responses
Denominator excludes: Respondents with children, but they do not have asthma
Denominator excludes: Respondents without children

Note: The tables for CASTHNO2 were produced using _CHILDWT as the weighting variable

Data was weighted using LANDWT.

DEMOGRAPHIC	RESPONDENT NUMBER		Yes			No			
GROUPS	TOTAL	WEIGHTED	N	%	C.I. (95%)	N	%	C.I. (95%)	
TOTAL	85	12,588	58	68.7	52.8-84.6	27	31.3	15.4-47.2	
Female	57	6,913	40	75.1	62.9-87.3	17	24.9	12.7-37.1	
Non-Native	81	12,089	56	69.2	52.7-85.7	25	30.8	14.3-47.3	

TABLE 5: DOES THE CHILD STILL HAVE ASTHMA? (CASTHNO2)

Denominator excludes: Respondents who are female or respondents with do not know/refused or with missing responses

Denominator excludes: Respondents with children, but they do not have asthma

Denominator excludes: Respondents without children

Note: The tables for CASTHNO2 were produced using _CHILDWT as the weighting variable

There is not enough data (less than 50 total observations) to print this table.

^{&#}x27;% = Percentage, C.I.(95%) = Confidence Interval (at 95 percent probability level). Percentages are weighted to population characteristics.

Rows with less than 50 observations have been suppressed.

TABLE 6: DOES THE CHILD STILL HAVE ASTHMA? (CASTHNO2)

Denominator excludes: Respondents who are male or respondents with do not know/refused or with missing responses

Denominator excludes: Respondents with children, but they do not have asthma

Denominator excludes: Respondents without children

Note: The tables for CASTHNO2 were produced using _CHILDWT as the weighting variable

Data was weighted using _LANDWT.

DEMOGRAPHIC	RESPONDENT NUMBER		Yes			No		
GROUPS	TOTAL	WEIGHTED	N	%	C.I. (95%)	N	%	C.I. (95%)
Female	57	6,913	40	75.1	62.8-87.4	17	24.9	12.6-37.2
Non-Native	54	6,560	39	77.1	64.8-89.4	15	22.9	10.6-35.2

BEHAVIORAL RISK FACTOR SURVEILLANCE SYSTEM NORTH DAKOTA MODULE QUESTIONS DATA REPORT, 2011

End of Module Data Report

93 of 93 July 26, 2012