

West Market Gram Positive Antibigram
% Susceptible

Organisms	Isolates	Ampicillin	Cefepime	Cefotaxime	Cefotaxime, Non-meningitis	Ceftriaxone	Ceftriaxone, Non-meningitis	Clindamycin	Daptomycin	Erythromycin	Gentamicin	Levofloxacin	Linezolid	Meropenem	Nitrofurantoin	Oxacillin	Penicillin	Penicillin, Oral Non-Meningitis	Penicillin, Non-Meningitis	Rifampin	Tetracycline	Trimethoprim/Sulfamethoxazole	Vancomycin
Enterococcus faecalis	377	99							86			84	99		100		98				32		100
Enterococcus faecalis VRE																							
Enterococcus faecium	20	45							100			41	95				45				66		100
Enterococcus faecium VRE																							
Staphylococcus aureus	766					99		82	100	72	98	91	99		100	99	0			99	95	98	99
Methicillin Resistant Staphylococcus aureus (MRSA)	279					97		74	100	15	98	40	100		100	0	0			98	91	97	100
Staphylococcus lugdunensis	77							80	100	79	100	100	100		100	97	0			100	94		100
Coag-Negative Staphylococcus species	50							66	100	58	100	75	100		100	74	0			100	90		100
Streptococcus pneumoniae	48			95	97		97	97		75		100		89			79	79	100		89	89	100
Streptococcus pyogenes (Group A)																							
Streptococcus agalactiae (Group B)	40		100	100			100			30		100	100				100						100
Beta-Streptococcus, Non-Group A or B																							
Streptococcus viridans group																							

Percentages derived from fewer than 10 isolates may not be statistically significant; not all isolates tested against every antibiotic.

Nitrofurantoin results apply to Urine isolates only.

Isolates identified as Staphylococcus aureus are considered to be methicillin susceptible (MSSA).

Antibiotic reporting noting additional non-meningitis context reflects isolate sensitivities from non-spinal-fluid source for Streptococcus pneumoniae isolates.

■ >= 90% of isolates susceptible
■ 61-89% of isolates susceptible
■ <= 60% of isolates susceptible

West Market Gram Negative Antibigram
% Susceptible

Organisms	Isolates	Amikacin	Ampicillin	Cefazolin	Cefazolin, Urine	Cefepime	Ceftazidime	Ceftazone	Ciprofloxacin	Gentamicin	Levofloxacin	Meropenem	Nitrofurantoin	Piperacillin/Tazobactam	Tobramycin	Trimethoprim/Sulfamethoxazole
Acinetobacter species	22	100	—	—	—	90	90	59	100	90	95	95	—	—	95	100
Citrobacter freundii/freundii complex	120	100	0	0	—	94	82	80	91	97	93	95	93	90	94	93
Klebsiella (Enterobacter) aerogenes	77	100	0	0	—	94	76	75	98	100	100	93	21	80	100	100
Enterobacter cloacae	4	100	0	0	—	100	75	50	100	100	100	100	0	75	100	75
Escherichia coli	3992	100	68	84	95	99	98	98	64	95	88	98	98	98	96	86
Escherichia coli ESBL	127	99	0	0	0	1	0	0	13	73	29	100	91	91	65	52
Klebsiella oxytoca	11	100	0	45	—	100	100	100	100	100	100	100	100	100	100	100
Klebsiella oxytoca ESBL	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Klebsiella pneumoniae	521	100	0	90	97	99	99	99	93	99	96	99	33	96	99	95
Klebsiella pneumoniae ESBL	16	100	0	0	6	0	0	0	6	56	6	100	43	81	53	31
Klebsiella pneumoniae/oxytoca ESBL	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Morganella morganii	41	100	0	0	—	100	97	95	78	85	78	97	0	97	92	78
Pantoea agglomerans	2	100	100	50	—	100	100	100	100	100	100	100	—	100	100	100
Proteus mirabilis	277	100	93	0	97	99	99	96	90	96	90	—	0	100	96	92
Proteus mirabilis ESBL	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Providencia species	17	100	0	0	—	94	88	94	70	93	70	100	0	94	93	100
Pseudomonas aeruginosa	256	98	—	—	—	94	92	—	83	98	84	91	—	90	98	—
Serratia marcescens	38	100	0	0	—	97	94	89	94	100	94	94	0	100	94	—
Stenotrophomonas maltophilia	25	—	—	—	—	—	44	—	—	—	88	—	—	—	—	100

Percentages derived from fewer than 10 isolates may not be statistically significant; not all isolates tested against every antibiotic.

Nitrofurantoin results apply to Urine isolates only.

■ >= 90% of isolates susceptible
■ 61-89% of isolates susceptible
■ <= 60% of isolates susceptible