



West Market Gram Positive Antibigram
% Susceptible

Organisms	Isolates	Ampicillin	Cefepime	Cefotaxime	Cefotaxime, Non-meningitis	Ceftazidime	Ceftazidime	Ceftazidime, Non-meningitis	Clindamycin	Daptomycin	Erythromycin	Gentamicin	Levofloxacin	Linezolid	Meropenem	Nitrofurantoin	Oxacillin	Penicillin	Penicillin, Oral Non-Meningitis	Penicillin, Non-Meningitis	Rifampin	Tetracycline	Trimethoprim/Sulfamethoxazole	Vancomycin
Enterococcus faecalis	426	99							90			88	98		99		99					32		100
Enterococcus faecalis VRE																		53						
Enterococcus faecium	26	53								100		52	92									73		100
Enterococcus faecium VRE	12	0								100		0	100					0				25		0
Staphylococcus aureus	759					99			82	100	71	98	91	99		100	99	0			99	93	97	99
Methicillin Resistant Staphylococcus aureus (MRSA)	265					96			77	100	15	98	37	100		100	0				99	91	95	100
Staphylococcus lugdunensis	82								79	100	81	98	100	100		100	96	0			98	93		100
Coag-Negative Staphylococcus species	61								61	100	53	93	70	100		100	60	0			100	93		100
Streptococcus pneumoniae	72			91	96		94	98			66				83		70	70	98			86	81	100
Streptococcus pyogenes (Group A)																								
Streptococcus agalactiae (Group B)	37		100	100			100		52		41		100	100			100							100
Beta-Streptococcus, Non-Group A or B																								
Streptococcus viridans group																								

Percentages derived from fewer than 10 isolates may not be statistically significant; not all isolates tested against every antibiotic.

Nitrofurantoin results apply to Urine isolates only.

Isolates identified as Staphylococcus aureus are considered to be methicillin susceptible (MSSA).

Antibiotic reporting noting additional non-meningitis context reflects isolate sensitivities from non-spinal-fluid source for Streptococcus pneumoniae isolates.

■ >= 90% of isolates susceptible
■ 61-89% of isolates susceptible
■ <= 60% of isolates susceptible

West Market Gram Negative Antibigram
% Susceptible

Organisms	Isolates	Amikacin	Ampicillin	Cefazolin	Cefazolin, Urine	Cefepime	Ceftazidime	Ceftazone	Ciprofloxacin	Gentamicin	Levofloxacin	Meropenem	Nitrofurantoin	Piperacillin/Tazobactam	Tobramycin	Trimethoprim/Sulfamethoxazole
Acinetobacter species	24	100	—	—	—	87	87	62	91	87	91	100	—	—	87	100
Citrobacter freundii/freundii complex	122	100	0	0	—	96	81	81	91	98	95	99	92	86	97	94
Klebsiella (Enterobacter) aerogenes	87	100	0	0	—	96	76	77	98	100	98	95	16	81	100	98
Enterobacter cloacae	3	100	0	0	—	100	66	100	100	100	100	100	100	100	100	100
Escherichia coli	4196	99	69	83	95	99	98	98	85	95	89	98	98	98	96	86
Escherichia coli ESBL	156	100	0	0	0	0	0	0	8	78	27	99	92	92	70	51
Klebsiella oxytoca	9	100	0	60	—	100	77	100	100	100	100	100	100	100	100	100
Klebsiella oxytoca ESBL	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Klebsiella pneumoniae	573	99	0	91	97	99	98	98	94	99	96	99	36	96	98	95
Klebsiella pneumoniae ESBL	20	100	0	0	0	0	0	0	5	60	10	100	30	65	60	20
Klebsiella pneumoniae/oxytoca ESBL	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Morganella morganii	42	100	0	0	—	97	92	95	85	95	88	95	0	100	97	90
Pantoea agglomerans	2	100	50	0	—	100	100	100	100	100	100	100	—	100	100	100
Proteus mirabilis	322	100	91	0	93	99	98	96	90	94	91	—	0	99	92	91
Proteus mirabilis ESBL	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Providencia species	22	100	0	0	—	100	95	95	77	95	77	100	0	95	95	100
Pseudomonas aeruginosa	259	98	—	—	—	93	92	100	81	98	82	91	—	89	98	—
Serratia marcescens	37	100	0	0	—	97	97	91	91	100	91	94	0	97	88	—
Stenotrophomonas maltophilia	20	—	—	—	—	—	35	—	—	—	80	—	—	—	—	95

Percentages derived from fewer than 10 isolates may not be statistically significant; not all isolates tested against every antibiotic.

Nitrofurantoin results apply to Urine isolates only.

■ >= 90% of isolates susceptible
■ 61-89% of isolates susceptible
■ <= 60% of isolates susceptible