

Transition to Tenecteplase Checklist

- ❑ Involve all stakeholders. This should include pharmacy, providers, stroke program leadership, and administration
- ❑ Determine if this is a feasible change for your facility--the decision to switch is determined by your facility and providers*
- ❑ Consider following the State recommendations on dosing of Tenecteplase in stroke
- ❑ Develop a Tenecteplase order set
- ❑ Update all current stroke protocols and resources
- ❑ Develop education/competency check off for nursing, pharmacy, and providers
- ❑ Develop process to safely differentiate stroke and MI Tenecteplase dosing
- ❑ Set a timeline (Do not be afraid to extend timeline to ensure safe transition)
- ❑ Develop education for patients

Transition Tips

- ✓ Consider making a full transition to Tenecteplase rather than alternating between Tenecteplase and Alteplase
- ✓ Medication safety is the priority--utilize Tenecteplase rather than TNK or TNKase to avoid confusion with tPA or TXA
- ✓ Do not underestimate the number of places Alteplase verbiage can be found
- ✓ Share updated acute stroke order set with telemedicine services
- ✓ Update dot phrases and downtime forms with new order sets and protocols
- ✓ Be mindful that Tenecteplase cannot be exchanged if premixed and not administered

*The transition to Tenecteplase is a facility specific choice. The North Dakota Stroke Task Force does not endorse the use of one thrombolytic over the other.

1/2023

STROKE LITERATURE

- [Guidelines for the Early Management of Patients with Acute Ischemic Stroke: 2019 Update to the 2018 Guidelines for the Early Management of Acute Ischemic Stroke](#)
- [Tenecteplase Thrombolysis for Acute Ischemic Stroke](#)

DOSING	INCLUSION/EXCLUSION CRITERIA
<p>Tenecteplase dose for STROKE is 0.25mg/kg (actual body weight) with a MAXIMUM DOSE of 25 mg</p> <ul style="list-style-type: none"> • IV push over five seconds • Not compatible with any dextrose containing solutions • Flush IV with Normal Saline before and after Tenecteplase administration • You will never administer a full vial to treat stroke • The stroke dosage is NOT listed on the box 	<p>Please refer to the ND Inclusion and Exclusion Criteria for IV Thrombolytic Treatment of Ischemic Stroke</p>
COMPLICATIONS	
	<ul style="list-style-type: none"> • Oral Angioedema • Hemorrhagic Transformation • Systemic Bleeding

REVERSAL

If the patient experiences hemorrhage post-Tenecteplase administration, follow the Neurocritical Care Society and the Society of Critical Care Medicine Guidelines for reversal of thrombolytics:

- Cryoprecipitate 10 units IV

If Cryoprecipitate is contraindicated or unavailable, administer:

- Tranexamic acid 10 to 15 mg/kg IV push over 20 minutes

OR

- Aminocaproic acid 4 to 5 mg IV

MONITORING

-Vital signs and neuro checks every 15 minutes x 2 hours, every 30 minutes x 6 hours, then hourly until 24 hours after treatment

-If blood pressure is greater than 180/105, notify provider

-Repeat head CT if neuro status declines

-NIHSS – Post administration and with any neuro changes

-No anticoagulant/antiplatelet for 24 hours

-Monitor for complications