



West Market Gram Negative Antibigram
% Susceptible

Organisms	Isolates	Amikacin	Ampicillin	Ampicillin/Sulbactam	Cefepime	Ceftazidime	Ceftriaxone	Cephalothin	Ciprofloxacin	Gentamicin	Levofloxacin	Meropenem	Nitrofurantoin	Piperacillin/Tazobactam	Tetracycline	Tobramycin	Trimethoprim/Sulfamethoxazole
Acinetobacter species	25	100	0	—	100	88	44	—	100	100	100	96	0	100	—	100	88
Citrobacter freundii/freundii complex	107	100	0	—	97	74	74	—	83	96	86	98	93	81	—	96	86
Klebsiella (Enterobacter) aerogenes	64	100	0	—	96	80	79	—	96	98	96	98	15	84	—	98	100
Enterobacter cloacae	2	100	0	—	50	0	0	—	100	100	100	100	—	0	—	100	100
Escherichia coli	3716	100	68	—	99	98	98	—	84	95	88	98	98	98	—	96	86
Escherichia coli ESBL	107	99	0	—	10	0	0	—	10	75	25	99	93	92	—	73	39
Klebsiella oxytoca	1	100	0	—	100	100	100	—	100	100	100	100	—	100	—	100	100
Klebsiella oxytoca ESBL	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Klebsiella pneumoniae	511	100	0	—	99	98	99	—	93	99	95	98	35	98	—	99	96
Klebsiella pneumoniae ESBL	7	100	0	—	14	0	0	—	28	85	28	85	14	71	—	42	28
Klebsiella pneumoniae/oxytoca ESBL	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Morganella morganii	23	100	0	—	100	81	86	—	91	95	91	86	0	86	—	95	91
Pantoea agglomerans	8	100	62	—	75	87	87	—	100	100	100	75	100	87	—	87	100
Proteus mirabilis	235	100	93	—	99	98	98	—	86	97	87	—	0	99	—	96	95
Proteus mirabilis ESBL	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Providencia species	13	92	0	—	100	84	100	—	69	84	61	92	0	100	—	84	100
Pseudomonas aeruginosa	264	99	—	—	98	93	—	—	84	97	84	91	—	92	—	97	—
Serratia marcescens	54	100	0	—	98	92	83	—	87	100	88	100	0	92	—	84	—
Stenotrophomonas maltophilia	26	—	—	—	—	46	—	—	—	—	84	—	—	—	—	—	100

Percentages derived from fewer than 10 isolates may not be statistically significant; not all isolates tested against every antibiotic.

Nitrofurantoin results apply to Urine isolates only.

>= 90% of isolates susceptible
 61-89% of isolates susceptible
 <= 60% of isolates susceptible

West Market Gram Positive Antibigram
% Susceptible

Organisms	Isolates	Amoxicillin/Clavulanate	Ampicillin	Ampicillin/Sulbactam	Azithromycin	Cefaclor	Cefazolin	Cefepime	Cefotaxime	Cefotaxime, Non-meningitis	Ceftriaxone	Ceftriaxone, Non-meningitis	Cefuroxime	Ciprofloxacin	Clindamycin	Daptomycin	Erythromycin	Gatifloxacin	Gentamicin	Levofloxacin	Linezolid	Meropenem	Nitrofurantoin	Oxacillin	Penicillin	Penicillin, Oral Non-Meningitis	Penicillin, Non-Meningitis	Rifampin	Tetracycline	Trimethoprim/Sulfamethoxazole	Vancomycin
Enterococcus faecalis	313		99													80				86	99		99		98				30		99
Enterococcus faecalis VRE	—																														
Enterococcus faecium	18		61																	76	100				58				78		100
Enterococcus faecium VRE	10		0													100				0	100				0			14		0	
Staphylococcus aureus	760			100			100				99				82	100	73		99	91	100		100	100	0			99	94	98	100
Methicillin Resistant Staphylococcus aureus	295										97				71	100	20		96	48	100		100	0	0			100	89	97	100
Staphylococcus lugdunensis	62														82	100	86		100	100	100		100	93	0			96	95		100
Coag-Negative Staphylococcus species	59														62	100	53		94	78	98		100	62	0			98	93	100	100
Streptococcus pneumoniae	50								98	100		100	100				72					90			82	82	98		94	78	100
Streptococcus pyogenes (Group A)	—																														
Streptococcus agalactiae (Group B)	48							100	100						52		41			100	100				100						100
Beta-Streptococcus, Non-Group A or B	—																														
Streptococcus viridians group	—																														

Percentages derived from fewer than 10 isolates may not be statistically significant; not all isolates tested against every antibiotic.

Nitrofurantoin results apply to Urine isolates only.

Isolates identified as Staphylococcus aureus are considered to be methicillin susceptible (MSSA).

Antibiotic reporting noting additional non-meningitis context reflects isolate sensitivities from non-spinal-fluid source for Streptococcus pneumoniae isolates.

■■■■■ >= 90% of isolates susceptible
■■■■■ 61-89% of isolates susceptible
■■■■■ <= 60% of isolates susceptible