

## Rubella (German Measles)

Rubella, commonly called German measles, is a **vaccine-preventable** rash illness caused by the rubella virus. It is usually a mild disease but can be severe in some cases. **Rubella during pregnancy (congenital rubella) can cause severe complications, such as birth defects, miscarriage or fetal death.**

## Transmission

Rubella is spread **person to person** by droplets spread while sneezing or coughing. A person is infectious with rubella from one week before and until one week after onset of the rubella rash. A pregnant person may transmit rubella to their fetus.

## Symptoms

In children, rubella is usually mild, with few noticeable symptoms. For children who do have symptoms, a widespread **red rash** is typically the first sign. The rash generally first appears on the face and then spreads to the rest of the body, lasting about three days. Other nonspecific symptoms that may appear prior to the rash may include cold-like symptoms, fever, pink eye.

Most adults who get rubella usually have a mild illness, with low-grade fever, sore throat, and a rash that starts on the face and spreads to the rest of the body.

Many people with rubella will experience few to no symptoms.

## Diagnosis

A health care provider can diagnose rubella. A laboratory test is needed to confirm the diagnosis. Because rubella disease is eliminated in the United States, a person likely would have travelled recently to another country or been exposed to someone who recently travelled. Diagnosis in the U.S. requires consultation with the NDHHS and CDC.

## Treatment

There is no treatment for rubella. Health care providers will recommend supportive care if necessary.

## Prevention

All children between 12 and 15 months of age should be vaccinated with the first dose of MMR (measles, mumps and rubella vaccine). A booster dose of MMR is recommended at 4 to 6 years of age. Two doses of mumps-containing vaccine are about 88 percent effective in preventing illness. **Children are required to submit documentation of age-appropriate immunization against mumps to attend early child care facilities, schools and colleges in North Dakota.**

- The immunization status of all children and staff at schools or early childhood facilities should be reviewed.
- During mumps outbreaks, exposed children and staff who have not been immunized should be excluded until they are vaccinated, or, if they refuse vaccination, they should be quarantine for an appropriate amount of time following exposure.
- During rubella outbreaks, the NDHHS may recommend that certain individuals receive a third dose of MMR vaccine.

## Exclusion Guidance

Rubella is a highly contagious disease and poses a serious risk to pregnant people. People with rubella must be excluded from work, school, or child care until no longer infectious. Outbreak situations may also require exclusion of unvaccinated persons, as well.

For additional information about rubella, contact the North Dakota Department of Health and Human Services' Public Health Division at 800.472.2180.

 **Report Immediately: 800.472.2180 or 701.328.2378**

**This disease is a reportable condition. As mandated by North Dakota law, any incidence of this disease in humans shall be reported to the North Dakota Department of Health and Human Services.**

### Resources:

1. Centers for Disease Control and Prevention. (2020, December 31). *CDC Rubella (German measles)*. Centers for Disease Control and Prevention. Retrieved June 28, 2023, <https://www.cdc.gov/rubella/about/transmission.html>.
2. Kimberlin, D. W., Barnett, E. D., Lynfield, R., Sawyer, M. H. (2021) Red Book: 2021-2024 Report of the Committee on Infectious Diseases. 32<sup>nd</sup> ed. American Academy of Pediatrics. [Management and Prevention of Infectious Diseases; Rubella] [pages 648-655].