

Orthopoxvirus (JYNNEOS™) Vaccine Screening Form

For patients: The following questions will help us determine which vaccines you may be given today. If you answer “yes” to any question, it does not necessarily mean you should not be vaccinated. It just means additional questions must be asked. If a question is not clear, please ask your healthcare provider to explain it.

Patient’s name (last, first, middle): _____

Date of birth: _____

Please complete the following, the person giving you the vaccine may ask for more information based on your answers:

Yes	No	Unknown	Question
Yes	No	Unknown	In the past 14 days, have you been exposed to a person who has been notified by public health they have an orthopox infection (i.e., monkeypox)?
Yes	No	Unknown	Do you identify as gay, bisexual, transgender or male who engages in intimate contact with men?
Yes	No	Unknown	Are you 18 years or older?
Yes	No	Unknown	Have you had a severe allergic reaction (e.g., anaphylaxis) after a previous dose of the JYNNEOS™ vaccine?
Yes	No	Unknown	Have you had a severe allergic reaction (e.g., anaphylaxis) following gentamicin or ciprofloxacin?
Yes	No	Unknown	Have you had a severe allergic reaction (e.g., anaphylaxis) to chicken or egg protein and are currently avoiding exposure to all chicken or egg products?
Yes	No	Unknown	Have you received COVID-19 vaccine within the past 4 weeks?
Yes	No	Unknown	Have you previously been diagnosed with monkeypox?
Yes	No	Unknown	Are you feeling sick today?
Yes	No	Unknown	Are you pregnant or breastfeeding?
Yes	No	Unknown	Do you have a history of keloid scar formation?
Yes	No	Unknown	Do you work in a laboratory performing diagnostic testing for orthopoxviruses?
Yes	No	Unknown	Are you a health care provider that is indicated by public health officials to receive JYNNEOS™ vaccine as pre-exposure prophylaxis?

Question	Answer	Special Considerations
In the past 14 days, have you been exposed to a person who has been notified by public health they have an orthopox infection (i.e., monkeypox)?	Yes	Administer age appropriate Jynneos™ vaccine following guidance found in interim clinical considerations .
Do you identify as gay, bisexual, transgender or male who engages in intimate contact with men?	Yes	Administer age appropriate Jynneos™ vaccine following guidance found in interim clinical considerations .
	No	Do not administer Jynneos™ vaccine unless otherwise indicated for Pre-exposure Prophylaxis.
Are you 18 years or older?	Yes	Administer Jynneos™ vaccine intradermally unless otherwise indicated.
	No	Administer Jynneos™ vaccine subcutaneously.
Have you had a severe allergic reaction (e.g., anaphylaxis) after a previous dose of the JYNNEOS™ vaccine?	Yes	Do not vaccinate. Referral to an allergist-immunologist should be considered to assess the risks versus benefits of administering a dose.
Have you had a severe allergic reaction (e.g., anaphylaxis) following gentamicin or ciprofloxacin?	Yes	Discuss risks and benefits with potential recipients. They may be vaccinated with a 30-minute observation period. Alternatively, vaccination can be delayed until an allergist-immunologist is consulted, but the impact of delaying vaccination should be considered.
Have you had a severe allergic reaction (e.g., anaphylaxis) to chicken or egg protein and are currently avoiding exposure to all chicken or egg products?	Yes	Discuss risks and benefits with potential recipients. They may be vaccinated with a 30-minute observation period. Alternatively, vaccination can be delayed until an allergist-immunologist is consulted, but the impact of delaying vaccination should be considered.
Have you received COVID-19 vaccine within the past 4 weeks?	Yes	If an orthopoxvirus vaccine is offered for prophylaxis in the setting of a monkeypox outbreak, orthopoxvirus vaccination should not be delayed because of recent receipt of a Moderna, Novavax, or Pfizer-BioNTech COVID-19 vaccine; no minimum interval between COVID-19 vaccination with these vaccines and orthopoxvirus vaccination is necessary. People, particularly adolescent or young adult males, might consider waiting 4 weeks after orthopoxvirus vaccination before receiving a Moderna, Novavax, or Pfizer-BioNTech COVID-19 vaccine, because of the observed risk for myocarditis and/or pericarditis after receipt of ACAM2000 orthopoxvirus vaccine and mRNA (i.e., Moderna and Pfizer-BioNTech) and Novavax COVID-19 vaccines and the unknown risk for myocarditis and/or pericarditis after JYNNEOS.

Question	Answer	Special Considerations
Have you previously been diagnosed with monkeypox?	Yes	<p>A person who would be eligible for vaccination but has been diagnosed with monkeypox during this outbreak, which started in the United States on May 17, 2022, is not recommended to be vaccinated at this time.</p> <p>An immunocompromised person who is diagnosed with monkeypox after their first dose of JYNNEOS may be eligible to receive the second dose of JYNNEOS on a case-by-case shared decision-making basis based on the clinical judgment of the healthcare provider.</p>
Are you feeling sick today?	Yes	<p>If moderate or severe acute illness, with or without fever consider deferring vaccination until the acute illness has improved.</p> <p>People presenting with minor illnesses, such as a cold, may be vaccinated.</p>
Are you pregnant or breastfeeding?	Yes	<p>The safety and efficacy of Jynneos™ has not been evaluated in people who are breastfeeding or in young children. It is not known whether JYNNEOS is excreted in human milk. Data are not available to assess the impact of Jynneos™ on milk production or the safety of breast milk for children from persons vaccinated with Jynneos™. However, Jynneos™ vaccine is replication-deficient and therefore should not present a risk of transmission to breastfed infants.</p> <p>Jynneos™ can be offered to people who are pregnant or breastfeeding who are otherwise eligible. The risks and benefits of JYNNEOS should be discussed with the patient using shared decision-making.</p>
Do you have a history of keloid scar formation?	Yes	Administer Jynneos™ vaccine subcutaneously regardless of age or vaccine availability.
Do you work in a laboratory performing diagnostic testing for orthopoxviruses?	Yes	Administer Jynneos™ vaccine intradermally as appropriate.
Are you a health care provider that is indicated by public health officials to receive JYNNEOS™ vaccine as pre-exposure prophylaxis?	Yes	Administer Jynneos™ vaccine intradermally as appropriate.