


Pulse Oximetry Screening Protocol for Critical Congenital Heart Disease (CCHD)

Check pulse ox in right hand and either foot in room air for newborns over 24 hours of age or shortly before discharge if less than 24 hours.



Tips for Pulse Oximetry Screeners

Where do I place the sensor?



Photo courtesy of Masimo

EITHER FOOT:

Wrap the sensor around the outer aspect of <u>either</u> foot. Place the light emitter on the top of the foot with the photodetector directly opposite of it. The tape/wrap should be securely wrapped around the foot.



Photo courtesy of Masimo

RIGHT HAND:

Wrap the sensor around the outer aspect of the <u>right</u> hand. Place the light emitter on the top of the right hand with the photodetector directly opposite of it. The tape/wrap should be securely wrapped around the hand.

What do I say to the family?

SCREEN:

We are going to screen your baby for critical congenital heart disease using a fast and painless test called pulse oximetry. The test will only take a few minutes, and it checks the oxygen level in your baby's blood.

REPEAT SCREEN:

Your baby's oxygen level was lower than we'd like when we performed the pulse oximetry screen, so we will recheck it again in about an hour.

This project is supported by the Health Resources and Services Administration (HRSA) of the U.S. Department of Health and Human Services (HHS) under grant number B04MC29321, Maternal and Child Health Services, total award amount for the period of October 1, 2016 through September 30, 2017 is \$1,725,639. This information or content and conclusions are those of the author and should not be construed as the official position or policy of, nor should any endorsements be inferred by HRSA, HHS or the U.S. Government.

DID NOT PASS:

Your baby's oxygen level was low when we performed the pulse oximetry screen. Your baby's care team needs to follow-up right away to find out why the oxygen level was low. Sometimes a low oxygen level is a sign of health problems. If there is a problem, it is best to figure it out before your baby goes home.

PASS:

Your baby had a normal oxygen level when we performed the pulse oximetry screen. No further testing is needed at this time. Since screening does not detect all heart defects, it is important for you to know what symptoms to look for in the future. Watch your baby's H.E.A.R.T.: Heart rate (too fast or too slow?); Energy (overly sleepy or fussy?); Appearance (pale or blue skin?); Respiration (breathing too fast or too slow?); Temperature (cold to the touch?).