

## Immunization Schedule Using Combination Vaccines: Pediarix®

		2 months	4 months	6 months	12 months	15-18 months	4-6 years	11-12 years	13-18 years
Combination Vaccines	Birth Dose of HBV	Pediarix® DTaP/IPV/HBV	Pediarix® DTaP/IPV/HBV	Pediarix® DTaP/IPV/HBV	Catch-up	Catch-up	DTaP/IPV MMRV		
DTaP					Catch-up	DTaP	Catch-up		
PCV		PCV	PCV	PCV	PCV	Catch-up	Catch-up	High Risk	High Risk
Rotavirus		Rotavirus	Rotavirus	Rotavirus <sup>1</sup>					
Hib		Hib	Hib	Hib <sup>2</sup>	Hib	Catch-up	Catch-up		
MMR					MMR	Catch-up	Catch-up	Catch-up	Catch-up
Varicella					Varicella	Catch-up	Catch-up	Catch-up	Catch-up
Hepatitis A					Hepatitis A	Hepatitis A	Catch-up	Catch-up	Catch-up
Meningococcal ACYW-135		High risk only	High risk only	High risk only	High risk only	High risk only	High risk only	MCV4	MCV4 booster (at age 16)
Meningococcal B								High risk only	Men B (2 doses at age 16)
HPV								HPV (2 dose series) <sup>3</sup>	Catch-up
Tdap								Tdap	Catch-up
Influenza				Influenza- some children may need 2 doses					
COVID-19				COVID-19					

1. If a child received Rotateq® or an unknown brand of rotavirus vaccine for the first or second dose in the series, three doses of rotavirus are needed.
2. If only using PedvaxHib®, a dose at 6 months is not necessary.
3. Those who initiate the HPV vaccine series before their 15<sup>th</sup> birthday and are immunocompetent only need two doses.

Contact the Immunization Unit:  
800.472.2180  
[www.hhs.nd.gov/immunizations](http://www.hhs.nd.gov/immunizations)