
## Pneumococcal Vaccine Recommendations: Adults ages 65+ years

The CDC recommends pneumococcal vaccination for adults 65 years and older who have no previously received a pneumococcal conjugate vaccine or whose previous vaccination history is unknown



## Pneumococcal Vaccine Recommendations: Adults ages 65+ years

The CDC recommends pneumococcal vaccination for adults 65 years and older who have no previously received a pneumococcal conjugate vaccine or whose previous vaccination history is unknown

## 1) Immunocompromising conditions include:

- Chronic renal failure
- Congenital or acquired asplenia
- Congenital or acquired immunodeficiency (including B-humoral) or T lymphocyte deficiency, complement deficiencies (particularly C1, C2, C3, and C4 deficiencies), and phagocytic disorders (excluding chronic granulomatous disease)
- Generalized malignancy
- HIV infection
- Hodgkin disease
- latrogenic immunosuppression (including diseases requiring treatment with immunosuppressive drugs, including long-term systemic corticosteroids and radiation therapy)
- Leukemia
- Lymphoma
- Multiple myeloma
- Nephrotic syndrome
- Sickle cell disease
- Solid organ transplants

## 2) Underlying medical conditions or other risk factors include:

- Cochlear implant
- Cerebrospinal fluid leak
- 3) If PPSV23 is administered, use shared clinical decision-making to decide whether to administer one dose of PCV20 at least 5 years after the last PPSV23 dose.



The PneumoRecs VaxAdvisor mobile app helps vaccination providers quickly and easily determine which pneumococcal vaccines a patient needs and when. The app incorporates recommendations for all ages so internists, family physicians, pediatricians, and pharmacists alike will find the tool beneficial. The app provides patient-specific guidance consistent with the immunization schedule recommended by the U.S. Advisory Committee on Immunization Practices (ACIP). <u>Download the App Today!</u>

