

Immunization Schedule Using Combination Vaccines: Vaxelis™

Health & Human Services

	Birth Dose of HBV	2 months	4 months	6 months	12 months	15-18 months	4-6 years	11-12 years	13-18 years
Combination Vaccines	Vaxelis™ DTaP/IPV/Hib/HBV	Vaxelis™ DTaP/IPV/Hib/HBV	Vaxelis™ DTaP/IPV/Hib/HBV	Catch-up	Pentacel® DTaP/IPV-Hib	DTaP/IPV MMRV			
PCV	PCV	PCV	PCV	PCV	Catch-up	Catch-up	High Risk	High Risk	
Rotavirus	Rotavirus	Rotavirus	Rotavirus ¹						
MMR				MMR	Catch-up	Catch-up	Catch-up	Catch-up	Catch-up
Varicella				Varicella	Catch-up	Catch-up	Catch-up	Catch-up	Catch-up
Hepatitis A				Hepatitis A	Hepatitis A	Catch-up	Catch-up	Catch-up	Catch-up
Meningococcal ACYW-135	High risk only	High risk only	High risk only	High risk only	High risk only	High risk only	High risk only	MCV4	MCV4 booster (at age 16)
Meningococcal B								High risk only	Men B (2 doses at age 16)
HPV								HPV (2 dose series) ²	Catch-up
Tdap								Tdap	Catch-up
Influenza			Influenza- some children may need 2 doses						
COVID-19			COVID-19						

1. If a child received Rotateq® or an unknown brand of rotavirus vaccine for the first or second dose in the series, three doses of rotavirus are needed.
2. Those who initiate the HPV vaccine series before their 15th birthday and are immunocompetent only need two doses.